

# PKN $\beta$ siRNA (m): sc-62823

## BACKGROUND

PKN $\beta$  (protein kinase PKN $\beta$ ), also known as PKN3, is a protein kinase-related molecule belonging to the AGC Serine/Threonine protein kinase family. It contains one protein kinase domain, three REM repeats and one AGC-kinase domain at its C-terminus. PKN $\beta$  is not expressed in normal adult tissues but is found in prostate tumors and various other cancer cell lines localizing to the nucleus and the perinuclear region of the cytoplasm. PKN $\beta$  may play a role in the invasiveness of malignant prostate cancer. This is suggested by the impaired growth and reduced metastases formation after knockdown of PKN $\beta$  expression in mouse prostate tumor cells. PKN $\beta$  expression and activity is regulated by PI 3-kinase. In humans, the phosphorylation of PKN $\beta$  at Thr 718 and Thr 860 is required for the activation of its kinase activity.

## REFERENCES

- Oishi, K., Mukai, H., Shibata, H., Takahashi, M. and Ono, Y. 1999. Identification and characterization of PKN $\beta$ , a novel isoform of protein kinase PKN: expression and arachidonic acid dependency are different from those of PKN $\alpha$ . *Biochem. Biophys. Res. Commun.* 261: 808-814.
- Shibata, H., Oishi, K., Yamagiwa, A., Matsumoto, M., Mukai, H. and Ono, Y. 2001. PKN $\beta$  interacts with the SH3 domains of Graf and a novel Graf related protein, Graf2, which are GTPase activating proteins for Rho family. *J. Biochem.* 130: 23-31.
- Oishi, K., Takahashi, M., Mukai, H., Banno, Y., Nakashima, S., Kanaho, Y., Nozawa, Y. and Ono, Y. 2001. PKN regulates phospholipase D1 through direct interaction. *J. Biol. Chem.* 276: 18096-18101.
- Online Mendelian Inheritance in Man, OMIM<sup>™</sup>. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 610714. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Leenders, F., Möpert, K., Schmiedeknecht, A., Santel, A., Czauderna, F., Aleku, M., Penschuck, S., Dames, S., Sternberger, M., Röhl, T., Wellmann, A., Arnold, W., Giese, K., Kaufmann, J. and Klippel, A. 2004. PKN3 is required for malignant prostate cell growth downstream of activated PI 3-kinase. *EMBO J.* 23: 3303-3313.
- Mukai, H. and Ono, Y. 2006. Purification and kinase assay of PKN. *Meth. Enzymol.* 406: 234-250.
- Wissing, J., Jänsch, L., Nimtz, M., Dieterich, G., Hornberger, R., Keri, G., Wehland, J. and Daub, H. 2007. Proteomics analysis of protein kinases by target class-selective prefractionation and tandem mass spectrometry. *Mol. Cell. Proteomics* 6: 537-547.

## CHROMOSOMAL LOCATION

Genetic locus: Pkn3 (mouse) mapping to 2 B.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## PRODUCT

PKN $\beta$  siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PKN $\beta$  shRNA Plasmid (m): sc-62823-SH and PKN $\beta$  shRNA (m) Lentiviral Particles: sc-62823-V as alternate gene silencing products.

For independent verification of PKN $\beta$  (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-62823A, sc-62823B and sc-62823C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

PKN $\beta$  siRNA (m) is recommended for the inhibition of PKN $\beta$  expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PKN $\beta$  gene expression knockdown using RT-PCR Primer: PKN $\beta$  (m)-PR: sc-62823-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.