

# T $\beta$ -10 siRNA (h): sc-63094

## BACKGROUND

Thymosin  $\beta$ -10 (T $\beta$ -10) is a member of the highly conserved  $\beta$ -thymosin family. It is a monomeric G-Actin sequestering protein of the cytoplasm that regulates Actin dynamics. T $\beta$ -10 consists of 43 amino acids and often forms  $\alpha$ -helical structures. T $\beta$ -10 has been shown to act as an Actin-mediated tumor suppressor. Overexpression of this protein inhibits endothelial cell proliferation, migration, invasion and tube formation. In human ovarian cancer cells, T $\beta$ -10 also increases apoptosis frequency. T $\beta$ -10 directly interacts with Ras, resulting in inhibition of the Ras downstream signaling pathways which, in turn, exhibits a negative effect on angiogenesis and tumor growth. More specifically, this inhibitive effect might be mediated by the downregulation of vascular endothelial growth factor (VEGF), VEGF receptor-1 (VEGFR-1) and Integrin  $\alpha$ V, which suggests a role for T $\beta$ -10 in anticancer therapy.

## REFERENCES

- Chiappetta, G., et al. 2004. Thymosin  $\beta$ -10 gene expression as a possible tool in diagnosis of thyroid neoplasias. *Oncol. Rep.* 12: 239-243.
- Alldinger, I., et al. 2005. Gene expression analysis of pancreatic cell lines reveals genes overexpressed in pancreatic cancer. *Pancreatology* 5: 370-379.
- Meeuwssen, S., et al. 2005. Cultured human adult microglia from different donors display stable cytokine, chemokine and growth factor gene profiles but respond differently to a pro-inflammatory stimulus. *Neuroimmunomodulation* 12: 235-245.
- Lee, S.H., et al. 2005. Thymosin  $\beta$ -10 inhibits angiogenesis and tumor growth by interfering with Ras function. *Cancer Res.* 65: 137-148.
- Rho, S.B., et al. 2005. The identification of apoptosis-related residues in human thymosin  $\beta$ -10 by mutational analysis and computational modeling. *J. Biol. Chem.* 280: 34003-34007.
- Huang, C.M., et al. 2006. *In vivo* detection of secreted proteins from wounded skin using capillary ultrafiltration probes and mass spectrometric proteomics. *Proteomics* 6: 5805-5814.
- Mu, H., et al. 2006. Thymosin  $\beta$ -10 inhibits cell migration and capillary-like tube formation of human coronary artery endothelial cells. *Cell Motil. Cytoskeleton* 63: 222-230.

## CHROMOSOMAL LOCATION

Genetic locus: TMSB10 (human) mapping to 2p11.2.

## PRODUCT

T $\beta$ -10 siRNA (h) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see T $\beta$ -10 shRNA Plasmid (h): sc-63094-SH and T $\beta$ -10 shRNA (h) Lentiviral Particles: sc-63094-V as alternate gene silencing products.

For independent verification of T $\beta$ -10 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-63094A and sc-63094B.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

T $\beta$ -10 siRNA (h) is recommended for the inhibition of T $\beta$ -10 expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

T $\beta$ -10 (D-6): sc-514309 is recommended as a control antibody for monitoring of T $\beta$ -10 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.