

# TIF1 $\gamma$ siRNA (h): sc-63127

## BACKGROUND

Transcriptional intermediary factor 1- $\alpha$  (TIF1 $\alpha$ ) mediates transcriptional events by interactions with the AF2 region of several nuclear receptors, such as the estrogen, retinoic acid, and vitamin D<sub>3</sub> receptors. TIF1 $\alpha$  localizes to nuclear bodies and is a member of the tripartite motif (TRIM) family. The TRIM motif includes three zinc-binding domains (RING, B-box type 1 and B-box type 2) and a coiled-coil region. TIF1 $\beta$  is also a member of the TRIM family that contains both a Cys/His PHD finger and bromodomain that form a cooperative unit required for transcriptional repression. TIF1 $\beta$  mediates transcriptional control by interaction with the Krüppel-associated box (KRAB) repression domain found in many transcription factors and by binding DNA via its zinc finger. TIF1 $\gamma$  has a similar structure to the previous two TRIM members, though it presents several functional differences. TIF1 $\gamma$  interacts with the Smad2/3 transcription factor in hematopoietic, mesenchymal, and epithelial cell types to mediate different transcriptional effects in response to TGF $\beta$ .

## REFERENCES

1. Friedman, J., et al. 1996. KAP-1, a novel corepressor for the highly conserved KRAB repression domain. *Genes Dev.* 10: 2067-2078.
2. Moosmann, P., et al. 1996. Transcriptional repression by RING finger protein TIF1 $\beta$  that interacts with the KRAB repressor domain of KOX1. *Nucleic Acids Res.* 24: 4859-4867.
3. Venturini, L., et al. 1999. TIF1 $\gamma$ , a novel member of the transcriptional intermediary factor 1 family. *Oncogene* 18: 1209-1217.

## CHROMOSOMAL LOCATION

Genetic locus: TRIM33 (human) mapping to 1p13.2.

## PRODUCT

TIF1 $\gamma$  siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TIF1 $\gamma$  shRNA Plasmid (h): sc-63127-SH and TIF1 $\gamma$  shRNA (h) Lentiviral Particles: sc-63127-V as alternate gene silencing products.

For independent verification of TIF1 $\gamma$  (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-63127A, sc-63127B and sc-63127C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## APPLICATIONS

TIF1 $\gamma$  siRNA (h) is recommended for the inhibition of TIF1 $\gamma$  expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

TIF1 $\gamma$  (XX-19): sc-101179 is recommended as a control antibody for monitoring of TIF1 $\gamma$  gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>TM</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TIF1 $\gamma$  gene expression knockdown using RT-PCR Primer: TIF1 $\gamma$  (h)-PR: sc-63127-PR (20  $\mu$ l, 557 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## SELECT PRODUCT CITATIONS

1. Lee, E.J., et al. 2020. Hepatic stellate cell-specific knockout of transcriptional intermediary factor 1 $\gamma$  aggravates liver fibrosis. *J. Exp. Med.* 217: e20190402.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.