



S-100A13 siRNA (m): sc-63356

BACKGROUND

The S-100 protein family consists of a group of calcium-binding proteins, that exhibit cell and tissue-specific expression. The expression levels of its members differ in various pathological conditions. The extracellular functions of the S-100 family may include the ability to enhance neurite outgrowth, involvement in inflammation and motility of tumor cells. S-100A13 is a 98 amino acid protein with two EF-hand calcium-binding domains. High levels of S-100A13 mRNA are present in skeletal muscle, heart, kidney, ovary, small intestine and pancreas. S-100A13 translocates in response to elevated intracellular calcium levels induced by Angiotensin II. S-100A13 co-localizes with S-100A1 on human chromosome 1q21, the site where the majority of S-100 proteins cluster.

REFERENCES

1. Wicki, R., et al. 1996. Characterization of the human and mouse cDNAs coding for S-100A13, a new member of the S-100 protein family. *Biochem. Biophys. Res. Commun.* 227: 594-599.
2. Mouta Carreira, C., et al. 1998. S-100A13 is involved in the regulation of fibroblast growth factor-1 and p40 synaptotagmin-1 release *in vitro*. *J. Biol. Chem.* 273: 22224-22231.
3. Online Mendelian Inheritance in Man, OMIM™. 1998. Johns Hopkins University, Baltimore, MD. MIM Number: 601989. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Shishibori, T., et al. 1999. Three distinct anti-allergic drugs, amlexanox, cromolyn and tranilast, bind to S-100A12 and S-100A13 of the S-100 protein family. *Biochem. J.* 338: 583-589.
5. Hsieh, H.L., et al. 2002. S-100A13 and S-100A6 exhibit distinct translocation pathways in endothelial cells. *J. Cell Sci.* 115: 3149-3158.
6. Chan, W.Y., et al. 2003. Differential expression of S-100 proteins in the developing human hippocampus and temporal cortex. *Microsc. Res. Tech.* 60: 600-613.
7. Hsieh, H.L., et al. 2004. S-100 protein translocation in response to extracellular S-100 is mediated by receptor for advanced glycation endproducts in human endothelial cells. *Biochem. Biophys. Res. Commun.* 16: 949-959.

CHROMOSOMAL LOCATION

Genetic locus: S100a13 (mouse) mapping to 3 F1.

PRODUCT

S-100A13 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see S-100A13 shRNA Plasmid (m): sc-63356-SH and S-100A13 shRNA (m) Lentiviral Particles: sc-63356-V as alternate gene silencing products.

For independent verification of S-100A13 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-63356A, sc-63356B and sc-63356C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

S-100A13 siRNA (m) is recommended for the inhibition of S-100A13 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor S-100A13 gene expression knockdown using RT-PCR Primer: S-100A13 (m)-PR: sc-63356-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.