

B lymphocytes (Lym-1): sc-69676

BACKGROUND

B lymphocytes are cells that play a large role in the humoral immune response, as opposed to the cell-mediated immune response that is governed by T cells. The principal function of B lymphocytes is to make antibodies against soluble antigens although, they do not produce antibodies until they become fully activated. B lymphocytes have unique receptor proteins (referred to as the B cell receptors (BCRs)) on their surfaces that will bind to one particular antigen. BCRs are composed of membrane-bound immunoglobulin, and allow for the distinction of B lymphocytes from other types of lymphocytes, as well as being the principal proteins involved in B lymphocyte activation. The human body makes millions of different types of B lymphocytes each day that circulate in the blood and lymph nodes.

REFERENCES

1. Armitage, R.J., Rowe, D.J. and Beverly, P.C. 1988. A new antigen identified by the monoclonal antibody UCHB 1 delivers a costimulatory signal to a subset of human B cells. *Eur. J. Immunol.* 18: 67-76.
2. Monroe, J.G., Bannish, G., Fuentes-Panana, E.M., King, L.B., Sandel, P.C., Chung, J. and Sater, R. 2003. Positive and negative selection during B lymphocyte development. *Immunol. Res.* 27: 427-442.
3. Donahue, A.C. and Fruman, D.A. 2004. PI3K signaling controls cell fate at many points in B lymphocyte development and activation. *Semin. Cell Dev. Biol.* 15: 183-197.
4. Dunn-Walters, D.K., Edelman, H. and Mehr, R. 2004. Immune system learning and memory quantified by graphical analysis of B lymphocyte phylogenetic trees. *Biosystems* 76: 141-155.
5. Zouali, M. and Sarmay, G. 2004. B lymphocyte signaling pathways in systemic autoimmunity: implications for pathogenesis and treatment. *Arthritis Rheum.* 50: 2730-2741.
6. Rickert, R.C. 2005. Regulation of B lymphocyte activation by complement C3 and the B cell coreceptor complex. *Curr. Opin. Immunol.* 17: 237-243.
7. Skok, M., Grailhe, R. and Changeux, J.P. 2005. Nicotinic receptors regulate B lymphocyte activation and immune response. *Eur. J. Pharmacol.* 517: 246-251.
8. Tedder, T.F., Poe, J.C. and Haas, K.M. 2005. CD22: a multifunctional receptor that regulates B lymphocyte survival and signal transduction. *Adv. Immunol.* 88: 1-50.
9. Titanji, K., Chiodi, F., Bellocco, R., Schepis, D., Osorio, L., Tassandin, C., Tambussi, G., Grutzmeier, S., Lopalco, L. and De Milito, A. 2005. Primary HIV-1 infection sets the stage for important B lymphocyte dysfunctions. *AIDS* 19: 1947-1955.

CHROMOSOMAL LOCATION

Genetic locus: CD83 (human) mapping to 6p23.

SOURCE

B lymphocytes (Lym-1) is a mouse monoclonal antibody raised against Burkitt's lymphoma cell line Raji of human origin.

PRODUCT

Each vial contains 200 µg IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

B lymphocytes (Lym-1) is available conjugated to either phycoerythrin (sc-69676 PE) or fluorescein (sc-69676 FITC), 200 µg/ml, for IF, IHC(P) and FCM.

APPLICATIONS

B lymphocytes (Lym-1) is recommended for detection of HLA-DR10 and B cell lymphomas of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and flow cytometry (1 µg per 1 x 10⁶ cells).


Molecular Weight of B lymphocytes: 36 kDa.

Positive Controls: BYDP whole cell lysate: sc-364368, HeLa whole cell lysate: sc-2200 or CCRF-CEM cell lysate: sc-2225.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



B lymphocytes (Lym-1): sc-69676. Western blot analysis of B lymphocytes expression in HeLa (A), CCRF-CEM (B), IMR-32 (C), BJAB (D), WEHI-231 (E) and Ramos (F) whole cell lysates.



B lymphocytes (Lym-1): sc-69676. Western blot analysis of B lymphocytes expression in BYDP whole cell lysate.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.