



CD5 (HK231): sc-69734

BACKGROUND

CD5 (also designated Lyt-1) has been identified as a transmembrane glycoprotein that is expressed on 70% of normal peripheral blood lymphocytes and on virtually all T lymphocytes in thymus and peripheral blood. Activation of T cells through the T cell receptor (TCR) results in tyrosine phosphorylation of CD5, and the absence of CD5 renders T cells hyper-responsive to TCR-mediated activation. CD5 associates with the TCR/CD3-z chain and with the Src family kinase Lck p56. *In vitro* studies have shown a 10- to 15-fold increase in the kinase activity of Lck bound to CD5. The B cell antigen, CD72, serves as a receptor for CD5. The consequence of CD5 binding to its cognate receptor is still in question and likely plays a role in thymic selection.

REFERENCES

1. Davies, A.A., Ley, S.C. and Crumpton, M.J. 1992. CD5 is phosphorylated on tyrosine after stimulation of the T cell antigen receptor complex. *Proc. Natl. Acad. Sci. USA* 89: 6368-6372.
2. Jamin, C., Lamour, A., Pennec, Y.L., Hirn, M., Le Goff, P. and Youinou, P. 1993. Expression of CD5 and CD72 on T and B cell subsets in rheumatoid arthritis and Sjögren's syndrome. *Clin. Exp. Immunol.* 92: 245-250.
3. Jones, M., Cordell, J.L., Beyers, A.D., Tse, A.G. and Mason, D.Y. 1993. Detection of T and B cells in many animal species using cross-reactive anti-peptide antibodies. *J. Immunol.* 150: 5429-5435.
4. Lydyard, P.M., Lamour, A., MacKenzie, L.E., Jamin, C., Mageed, R.A. and Youinou, P. 1994. CD5+ B cells and the immune system. *Immunol. Lett.* 38: 159-166.
5. Raab, M., Yamamoto, M. and Rudd, C.E. 1994. The T cell antigen CD5 acts as a receptor and substrate for the protein-tyrosine kinase p56lck. *Mol. Cell. Biol.* 14: 2862-2870. PMID: 7513045
6. Plater-Zyberk, C., Taylor, P.C., Blaylock, M.G. and Maini, R.N. 1995. Anti-CD5 therapy decreases severity of established disease in collagen type II-induced arthritis in DBA/1 mice. *Clin. Exp. Immunol.* 98: 442-447.
7. Ekerfelt, C., Ernerudh, J., Solders, G. and Vrethem, M. 1995. CD5 expression on B cells may antibodies in patients with polyneuropathy associated with monoclonal gammopathy. *Clin. Exp. Immunol.* 101: 346-350.
8. Tarakhovsky, A., Kanner, S.B., Hombach, J., Ledbetter, J.A., Müller, W., Killeen, N. and Rajewsky, K. 1995. A role for CD5 in TCR-mediated signal transduction and thymocyte selection. *Science* 269: 535-537.
9. Koskinen, R., Göbel, T.W., Tregaskes, C.A., Young, J.R. and Vainio, O. 1998. The structure of avian CD5 implies a conserved function. *J. Immunol.* 160: 4943-4950.

CHROMOSOMAL LOCATION

Genetic locus: CD5 (human) mapping to 11q12.2.

SOURCE

CD5 (HK231) is a mouse monoclonal antibody raised against CLL cells of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

CD5 (HK231) is recommended for detection of CD5 of human origin by immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and flow cytometry (1 µg per 1 x 10⁶ cells).

Molecular Weight of CD5: 67 kDa.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.