SANTA CRUZ BIOTECHNOLOGY, INC.

c-Myc (1.N.2): sc-70469



BACKGROUND

c-Myc-, N-Myc- and L-Myc-encoded proteins function in cell proliferation, differentiation and neoplastic disease. Myc proteins are nuclear proteins with relatively short half lives. Amplification of the c-Myc gene has been found in several types of human tumors including lung, breast and colon carcinomas, while the N-Myc gene has been found amplified in neuroblastomas. The L-Myc gene has been reported to be amplified and expressed at high level in human small cell lung carcinomas. The presence of three sequence motifs in the c-Myc COOH terminus, including the leucine zipper, the helix-loophelix and a basic region, provided initial evidence for a sequence-specific binding function. A basic region helix-loop-helix leucine zipper motif (bHLH-Zip) protein, designated Max, specifically associates with c-Myc, N-Myc and L-Myc proteins. The Myc-Max complex binds to DNA in a sequence-specific manner under conditions where neither Max nor Myc exhibit appreciable binding. Max can also form heterodimers with at least two additional bHLH-Zip proteins, Mad and Mxi1, and Mad 1-Max dimers have been shown to repress transcription through interaction with mSin3.

CHROMOSOMAL LOCATION

Genetic locus: MYC (human) mapping to 8q24.21; Myc (mouse) mapping to 15 D1.

SOURCE

c-Myc (1.N.2) is a mouse monoclonal antibody raised against full length c-Myc of human origin.

PRODUCT

Each vial contains 200 $\mu g\, lgG_1$ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

c-Myc (1.N.2) is recommended for detection of c-Myc p67 and c-Myc tagged fusion proteins of mouse, rat, human and avian origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for c-Myc siRNA (h): sc-29226, c-Myc siRNA (m): sc-29227, c-Myc siRNA (r): sc-270149, c-Myc shRNA Plasmid (h): sc-29226-SH, c-Myc shRNA Plasmid (m): sc-29227-SH, c-Myc shRNA Plasmid (r): sc-270149-SH, c-Myc shRNA (h) Lentiviral Particles: sc-29226-V, c-Myc shRNA (m) Lentiviral Particles: sc-29227-V and c-Myc shRNA (r) Lentiviral Particles: sc-270149-V.

Molecular Weight of c-Myc: 67 kDa.

Positive Controls: c-Myc (m): 293T Lysate: sc-118892, HeLa whole cell lysate: sc-2200 or K-562 whole cell lysate: sc-2203.

STORAGE

Store at 4° C, **D0 NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

DATA





c-Myc (1.N.2): sc-70469. Western blot analysis of c-Myc expression in non-transfected: sc-117752 (**A**) and mouse c-Myc transfected: sc-118892 (**B**) 293T whole cell lysates.

c-Myc (1.N.2): sc-70469. Western blot analysis of c-Myc expression in HeLa whole cell lysate.

SELECT PRODUCT CITATIONS

- Li, X.W., et al. 2012. Inhibitory effect of calcitonin gene-related peptide on hypoxia-induced rat pulmonary artery smooth muscle cells proliferation: role of ERK1/2 and p27. Eur. J. Pharmacol. 679: 117-126.
- Chong, K.Y., et al. 2015. Wnt pathway activation and ABCB1 expression account for attenuation of proteasome inhibitor-mediated apoptosis in multidrug-resistant cancer cells. Cancer Biol. Ther. 16: 149-159.
- Rulina, A.V., et al. 2016. Distinct outcomes of CRL-Nedd8 pathway inhibition reveal cancer cell plasticity. Cell Death Dis. 7: e2505.
- Alaca, N., et al. 2017. Treatment with milk thistle extract (*Silybum marianum*), ursodeoxycholic acid, or their combination attenuates cholestatic liver injury in rats: role of the hepatic stem cells. Turk. J. Gastroenterol. 28: 476-484.
- Chen, X., et al. 2018. Plant flavonoid taxifolin inhibits the growth, migration and invasion of human osteosarcoma cells. Mol. Med. Rep. 17: 3239-3245.
- Bi, L., et al. 2019. The histone chaperone complex FACT promotes proliferative switch of G_n cancer cells. Int. J. Cancer 145: 164-178.
- 7. Zhang, T., et al. 2020. TBL1XR1 is involved in c-Met-mediated tumorigenesis of human nonsmall cell lung cancer. Cancer Gene Ther. 27: 136-146.
- Sarkar, R., et al. 2021. Japanese encephalitis virus capsid protein interacts with non-lipidated MAP1LC3 on replication membranes and lipid droplets. J. Gen. Virol. E-published.
- Liao, W.C., et al. 2022. Nuclear accumulation of KPNA2 impacts radioresistance through positive regulation of the PLSCR1-STAT1 loop in lung adenocarcinoma. Cancer Sci. 113: 205-220.



See **c-Myc (9E10): sc-40** for c-Myc antibody conjugates, including AC, HRP, FITC, PE, Alexa Fluor[®] 488, 546, 594, 647, 680 and 790.