


CD4 (3H739): sc-70674



The Power to Question

BACKGROUND

The T cell receptor (TCR) is a heterodimer composed of either α and β or γ and δ chains. CD3 chains and the CD4 or CD8 co-receptors are also required for efficient signal transduction through the TCR. The TCR is expressed on T helper and T cytotoxic cells that can be distinguished by their expression of CD4 and CD8; T helper cells express CD4 proteins and T cytotoxic cells display CD8. CD4 is also expressed on cortical cells, mature medullary thymocytes, microglial cells and dendritic cells. CD4 (also designated T4 and Leu 3) is a membrane glycoprotein that contains four extracellular immunoglobulin-like domains. The TCR in association with CD4 can bind class II MHC molecules presented by the antigen-presenting cells. The CD4 protein functions by increasing the avidity of the interaction between the TCR and an antigen-class II MHC complex. An additional role of CD4 is to function as a receptor for HIV.

REFERENCES

1. Maddon, P.J., et al. 1987. Structure and expression of human and mouse T4 genes. Proc. Natl. Acad. Sci. USA 84: 9155-9159.
2. Arthos, J., et al. 1989. Identification of the residues in human CD4 critical for the binding of HIV. Cell 57: 469-481.
3. Healey, D., et al. 1990. Novel anti-CD4 monoclonal antibodies separate human immunodeficiency virus infection and fusion of CD4 $^{+}$ cells from virus binding. J. Exp. Med. 172: 1233-1242.
4. Allison, J.P., et al. 1991. The immunobiology of T cells with invariant $\gamma\delta$ antigen receptors. Annu. Rev. Immunol. 9: 679-705.
5. Janeway, C.A., Jr. 1992. The T cell receptor as a multicomponent signalling machine: CD4/CD8 co-receptors and CD45 in T cell activation. Annu. Rev. Immunol. 10: 645-674.
6. Ehrlich, E.W., et al. 1993. T cell receptor interaction with peptide/major histocompatibility complex (MHC) and superantigen/MHC ligands is dominated by antigen. J. Exp. Med. 178: 713-722.
7. Julius, M., et al. 1993. Distinct roles for CD4 and CD8 as co-receptors in antigen receptor signalling. Immunol. Today 14: 177-183.
8. Vignali, D.A. 1994. The interaction between CD4 and MHC class II molecules and its effect on T cell function. Behring Inst. Mitt. 94: 133-147.
9. Schlossman, S., et al, eds. 1995. Leucocyte Typing V. Oxford: Oxford University Press.

CHROMOSOMAL LOCATION

Genetic locus: CD4 (human) mapping to 12p13.31.

SOURCE

CD4 (3H739) is a rat monoclonal antibody raised against rat cells transfected with CD4 of human origin.

PRODUCT

Each vial contains 200 μ g IgG $_1$ in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

CD4 (3H739) is recommended for detection of CD4 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and flow cytometry (1 μ g per 1 \times 10 6 cells).

Suitable for use as control antibody for CD4 siRNA (h): sc-29246, CD4 shRNA Plasmid (h): sc-29246-SH and CD4 shRNA (h) Lentiviral Particles: sc-29246-V.

Molecular Weight of CD4: 54 kDa.

Positive Controls: HL-60 whole cell lysate: sc-2209, CCRF-CEM cell lysate: sc-2225 or SUP-T1 whole cell lysate: sc-364796.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **CD4 (MT310): sc-19641** for CD4 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.