

# COL10A1 (0.N.232): sc-70869

## BACKGROUND

Collagen Type X is associated with hypertrophic chondrocytes of avian and mammalian growth plate tissues during the endochondral growth of long bones. It is a component of normal articular cartilage in adult human, growing pig and newborn rat, and it is also present during any disruption of normal metabolic status of articular cartilage that occur with osteoarthritis. Collagen Type X is composed of three identical  $\alpha$  1(X) chains, each containing a triple-helical region flanked by a short N-terminal sequence and a larger non-collagenous C-terminal (NC1) domain. Mutations in COL10A1, the gene encoding for Collagen Type X, are associated with metaphyseal dysplasia type Schmid (SMCD) and other related forms of metaphyseal dysplasia. SMCD is characterized by short-limbed dwarfism, an outward "flaring" of the lower rib cage, bowed legs, leg pain and a hip deformity that causes the thigh bone to angle toward the center of the body.

## REFERENCES

- Gadher, S.J., et al. 1988. Susceptibility of cartilage Collagens Type II, IX, X and XI to human synovial collagenase and neutrophil elastase. *Eur. J. Biochem.* 175: 1-7.
- Gadher, S.J., et al. 1989. Cleavage of Collagen Type X by human synovial collagenase and neutrophil elastase. *Matrix* 9: 109-115.
- Rucklidge, G.J., et al. 1996. Collagen Type X: a component of the surface of normal human, pig and rat articular cartilage. *Biochem. Biophys. Res. Commun.* 224: 297-302.
- Barber, R.E., et al. 1997. Partial characterization of the C-terminal non-collagenous domain (NC1) of Collagen Type X. *Biochem. J.* 320: 479-485.
- Jacenko, O. 2000. Genetic-engineered models of skeletal diseases. I. Collagen Type X. *Methods Mol. Biol.* 137: 471-490.
- Magee, C., et al. 2005. Sp3/Sp1 transcription activity regulates specific expression of Collagen Type X in hypertrophic chondrocytes. *J. Biol. Chem.* 280: 25331-25338.
- Chong, I.W., et al. 2006. Great potential of a panel of multiple hMTH1, SPD, ITGA11 and COL11A1 markers for diagnosis of patients with non-small cell lung cancer. *Oncol. Rep.* 16: 981-988.
- Goto, T., et al. 2006. Sp1 family of transcription factors regulates the human  $\alpha$  2 (XI) collagen gene (COL11A2) in Saos-2 osteoblastic cells. *J. Bone Miner. Res.* 21: 661-673.
- Sjöblom, T., et al. 2006. The consensus coding sequences of human breast and colorectal cancers. *Science* 314: 268-274.

## SOURCE

COL10A1 (0.N.232) is a mouse monoclonal antibody raised against full length native Collagen  $\alpha$ 1 Type X of avian origin.

## PRODUCT

Each vial contains 200  $\mu$ g IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS

COL10A1 (0.N.232) is recommended for detection of Collagen  $\alpha$ 1 Type X of avian origin by immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

Molecular Weight of COL10A1: 66 kDa.

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:  
 1) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

## STORAGE

Store at 4° C, **\*\*DO NOT FREEZE\*\***. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.