

Cytokeratin 10/13 (4A28): sc-70908

BACKGROUND

Cytokeratins comprise a diverse group of intermediate filament proteins (IFPs) that are expressed as pairs in both keratinized and non-keratinized epithelial tissue. Cytokeratins play a critical role in differentiation and tissue specialization and function to maintain the overall structural integrity of epithelial cells. Cytokeratins have been found to be useful markers of tissue differentiation which is directly applicable to the characterization of malignant tumors. For example, Cytokeratins 10 and 13 are expressed highly in a subset of squamous cell carcinomas while Cytokeratin 18 is expressed in a majority of adenocarcinomas and basal cell carcinomas.

REFERENCES

1. van der Velden, L.A., et al. 1993. Cytokeratin expression in normal and (pre) malignant head and neck epithelia: an overview. *Head Neck* 15: 133-146.
2. Silen, A., et al. 1994. Evaluation of a new tumor marker for Cytokeratin 8 and 18 fragments in healthy individuals and prostate cancer patients. *Prostate* 24: 326-332.
3. Marceau, N., et al. 1995. Cytokeratin expression, fibrillar organization and subtle function in liver cells. *Biochem. Cell Biol.* 73: 619-625.
4. Quillien, V., et al. 1995. Serum and tissue distribution of a fragment of Cytokeratin 19 (CYFRA 21-1) in lung cancer patients. *Anticancer Res.* 15: 2857-2863.
5. Silen, A., et al. 1995. A novel IRMA and ELISA for quantifying Cytokeratin 8 and 18 fragments in the sera of healthy individuals and cancer patients. *Scan. J. Clin. Lab. Invest.* 55: 153-161.

CHROMOSOMAL LOCATION

Genetic locus: KRT10/KRT13 (human) mapping to 17q21.2.

SOURCE

Cytokeratin 10/13 (4A28) is a mouse monoclonal antibody raised against cytoskeletal preparation extracted from human ectocervical epithelium.

PRODUCT

Each vial contains 200 µg IgG_{2a} kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Cytokeratin 10/13 (4A28) is recommended for detection of Cytokeratin 10 and Cytokeratin 13 of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500); specific for Cytokeratin 13 by IHC(P).


Molecular Weight of Cytokeratin 10/13: 50 kDa.

Positive Controls: A-431 whole cell lysate: sc-2201.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgGκ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



Cytokeratin 10/13 (4A28): sc-70908. Western blot analysis of Cytokeratin 10/13 expression in untreated (A) and chemically-treated (B, C) HeLa whole cell lysates. β-Actin (C4): sc-47778 used as loading control. Detection reagent used: m-IgG Fc BP-HRP: sc-525409.



Cytokeratin 10/13 (4A28): sc-70908. Western blot analysis of Cytokeratin 10/13 expression in A-431 whole cell lysate.

SELECT PRODUCT CITATIONS

1. Mallery, S.R., et al. 2014. Topical application of a mucoadhesive freeze-dried black raspberry gel induces clinical and histologic regression and reduces loss of heterozygosity events in premalignant oral intraepithelial lesions: results from a multicentered, placebo-controlled clinical trial. *Clin. Cancer Res.* 20: 1910-1924.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **Cytokeratin 10 (LH2): sc-53252** for Cytokeratin 10 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.