

# IKK $\gamma$ (1.T.26): sc-71331

## BACKGROUND

The transcription factor NF $\kappa$ B is retained in the cytoplasm in an inactive form by the inhibitory protein I $\kappa$ B. Activation of NF $\kappa$ B requires that I $\kappa$ B be phospho-rylated on specific serine residues, which results in targeted degradation of I $\kappa$ B. I $\kappa$ B kinase  $\alpha$  (IKK $\alpha$ ), previously designated CHUK, interacts with I $\kappa$ B $\alpha$  and specifically phosphorylates I $\kappa$ B $\alpha$  on Serine 32 and 36, the sites that trigger its degradation. IKK $\alpha$  appears to be critical for NF $\kappa$ B activation in response to proinflammatory cytokines. Phosphorylation of I $\kappa$ B by IKK $\alpha$  is stimulated by the NF $\kappa$ B inducing kinase (NIK), which itself is a central regulator for NF $\kappa$ B activation in response to TNF and IL-1. The functional IKK complex contains three subunits, IKK $\alpha$ , IKK $\beta$  and IKK $\gamma$  (also designated NEMO), and each appear to make essential contributions to I $\kappa$ B phosphorylation.

## REFERENCES

1. Verma, I.M., et al. 1995. Rel/NF $\kappa$ B/I $\kappa$ B family: intimate tales of association and dissociation. *Genes Dev.* 9: 2723-2735.
2. Thanos, D., et al. 1995. NF $\kappa$ B: a lesson in family values. *Cell* 80: 529-532.
3. Connelly, M.A. and Marcu, K.B. 1995. CHUK, a new member of the helix-loop-helix and leucine zipper families of interacting proteins, contains a serine-threonine kinase catalytic domain. *Cell. Mol. Biol. Res.* 41: 537-549.
4. Malinin, N.L., et al. 1997. MAP3K-related kinase involved in NF $\kappa$ B induction by TNF, CD95 and IL-1. *Nature* 385: 540-544.
5. DiDonato, J.A., et al. 1997. A cytokine-responsive I $\kappa$ B kinase that activates the transcription factor NF $\kappa$ B. *Nature* 388: 548-554.
6. Regnier, C.H., et al. 1997. Identification and characterization of an I $\kappa$ B kinase. *Cell* 90: 373-383.
7. Zandi, E., et al. 1997. The I $\kappa$ B kinase complex (IKK) contains two kinase subunits, IKK $\alpha$  and IKK $\beta$ , necessary for I $\kappa$ B phosphorylation and NF $\kappa$ B activation. *Cell* 91: 243-252.
8. Song, H.Y., et al. 1997. Tumor necrosis factor (TNF)-mediated kinase cascades: bifurcation of nuclear factor- $\kappa$ B and c-jun N-terminal kinase (JNK/SAPK) pathways at TNF receptor-associated factor 2. *Proc. Natl. Acad. Sci. USA* 94: 9792-9296.
9. Yamaoka, S., et al. 1998. Complementation cloning of NEMO, a component of the I $\kappa$ B kinase complex essential for NF $\kappa$ B activation. *Cell* 93: 1231-1240.

## CHROMOSOMAL LOCATION

Genetic locus: IKBKG (human) mapping to Xq28.

## SOURCE

IKK $\gamma$  (1.T.26) is a mouse monoclonal antibody raised against full length His-tagged IKK $\gamma$  of human origin.

## PRODUCT

Each vial contains 100  $\mu$ g IgG<sub>1</sub> in 1.0 ml PBS with < 0.1% sodium azide and 0.1% gelatin.

## APPLICATIONS


IKK $\gamma$  (1.T.26) is recommended for detection of IKK $\gamma$  of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunoprecipitation [1-2  $\mu$ g per 100-500  $\mu$ g of total protein (1 ml of cell lysate)].

Suitable for use as control antibody for IKK $\gamma$  siRNA (h): sc-29363, IKK $\gamma$  shRNA Plasmid (h): sc-29363-SH and IKK $\gamma$  shRNA (h) Lentiviral Particles: sc-29363-V.


Molecular Weight of IKK $\gamma$ : 48 kDa.

Positive Controls: IKK $\gamma$  (h): 293T Lysate: sc-116282, Jurkat whole cell lysate: sc-2204 or K-562 whole cell lysate: sc-2203.

## DATA



IKK $\gamma$  (1.T.26): sc-71331. Western blot analysis of IKK $\gamma$  expression in BJAB (A), Jurkat (B), K-562 (C), U-937 (D), SW-13 (E) and HeLa (F) whole cell lysates.



IKK $\gamma$  (1.T.26): sc-71331. Western blot analysis of IKK $\gamma$  expression in non-transfected 293T: sc-117752 (A), human IKK $\gamma$  transfected 293T: sc-116282 (B) and K-562 (C) whole cell lysates.

## SELECT PRODUCT CITATIONS

1. Kawamoto, M., et al. 2018. The novel selective pan-TRK inhibitor ONO-7579 exhibits antitumor efficacy against human gallbladder cancer *in vitro*. *Anticancer Res.* 38: 1979-1986.
2. Yilmaz, D.E., et al. 2023. NLRX1 ligand, docosahexaenoic acid, ameliorates LPS-induced inflammatory hyperalgesia by decreasing TRAF6/IKK/I $\kappa$ B $\alpha$ /NF $\kappa$ B signaling pathway activity. *Cell. Mol. Biol.* 69: 15-23.

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.



See **IKK $\gamma$  (F-10): sc-166398** for IKK $\gamma$  antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.