

p53 (3H2821): sc-71815



The Power to Question

BACKGROUND

p53, a DNA-binding, oligomerization domain- and transcription activation domain-containing tumor suppressor that upregulates growth arrest and apoptosis-related genes in response to stress signals, thereby influencing programmed cell death, cell differentiation and cell cycle control mechanisms. p53 localizes to the nucleus, yet can be chaperoned to the cytoplasm by the negative regulator MDM2, an E3 ubiquitin ligase that is upregulated in the presence of active p53, where MDM2 polyubiquitinates p53 for proteasome targeting. p53 fluctuates between latent and active (DNA-binding) conformations, and is differentially activated through post-translational modifications including phosphorylation and acetylation. Mutations in the DNA-binding domain (DBD) of p53, amino acids 110-286, can compromise energetically favorable association with *cis* elements and are implicated in several human cancers.

REFERENCES

1. Banks, L., et al. 1986. Isolation of human p53-specific monoclonal antibodies and their use in the studies of human p53 expression. *Eur. J. Biochem.* 159: 529-534.
2. Hupp, T.R., et al. 1992. Regulation of the specific DNA-binding function of p53. *Cell* 71: 875-886.
3. Levine, A.J. 1997. p53, the cellular gatekeeper for growth and division. *Cell* 88: 323-331.
4. Ashcroft, M., et al. 1999. Regulation of p53 stability. *Oncogene* 18: 7637-7643.
5. Soussi, T., et al. 2000. p53 website and analysis of p53 gene mutations in human cancer: forging a link between epidemiology and carcinogenesis. *Hum. Mutat.* 15: 105-113.
6. Chene, P. 2001. The role of tetramerization in p53 function. *Oncogene* 20: 2611-2617.

CHROMOSOMAL LOCATION

Genetic locus: Trp53 (mouse) mapping to 11 B3.

SOURCE

p53 (3H2821) is a mouse monoclonal antibody raised against SVA31E7 mouse SV40-transformed cell line.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

APPLICATIONS

p53 (3H2821) is recommended for detection of wild type p53 of mouse origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for p53 siRNA (m): sc-29436, p53 shRNA Plasmid (m): sc-29436-SH and p53 shRNA (m) Lentiviral Particles: sc-29436-V.


Molecular Weight of p53: 53 kDa.

Positive Controls: BC₃H1 cell lysate: sc-2299, WRL19L cell lysate: sc-3805 or mouse LacZ whole cell lysate: sc-364371.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



p53 (3H2821): sc-71815. Western blot analysis of p53 expression in mouse LacZ whole cell lysate.

SELECT PRODUCT CITATIONS

1. Koryllou, A., et al. 2009. Cell death induced by N-methyl-N-nitrosourea, a model S(N)1 methylating agent, in two lung cancer cell lines of human origin. *Apoptosis* 14: 1121-1133.
2. Abdelfadil, E., et al. 2013. Thymoquinone induces apoptosis in oral cancer cells through p38β inhibition. *Am. J. Chin. Med.* 41: 683-696.



See **p53 (A-1): sc-393031** for p53 antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.