

PML (5G191): sc-71910

BACKGROUND

The PML protein is a zinc finger transcription factor expressed as three major transcription products due to alternative splicing. The gene encoding human PML maps to chromosome 15q24.1. The t(15;17) (q22;q11.2-q12) chromosomal translocation of the retinoic acid receptor α (RAR α) gene occurs in virtually all cases of acute promyelocytic leukemia and results in the expression of a PML/RAR α chimeric protein. Myeloid precursor cells expressing the PML/RAR α chimera fail to differentiate and exhibit an increased growth rate consequent to diminished apoptosis. PML/RAR α transforms myeloid precursors by recruiting the nuclear co-repressor (N-CoR)-histone deacetylase complex that is essential to retinoic acid-dependent myeloid differentiation. PML/RAR α also recruits DNA methyltransferases in order to induce gene hypermethylation and silencing, which ultimately facilitates leukemogenesis.

REFERENCES

1. de The, H., et al. 1990. The t(15;17) translocation of acute promyelocytic leukaemia fuses the retinoic acid receptor α gene to a novel transcribed locus. *Nature* 347: 558-561.
2. Borrow, J., et al. 1990. Molecular analysis of acute promyelocytic leukemia breakpoint cluster region on chromosome 17. *Science* 249: 1577-1580.
3. Pandolfi, P.P., et al. 1991. Structure and origin of the acute promyelocytic leukemia myl/RAR α cDNA and characterization of its retinoid-binding and transactivation properties. *Oncogene* 6: 1285-1292.

CHROMOSOMAL LOCATION

Genetic locus: PML (human) mapping to 15q24.1.

SOURCE

PML (5G191) is a mouse monoclonal antibody N-terminal epitope corresponding to residues 37-51 of PML of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

PML (5G191) is recommended for detection of all isoforms of PML of human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for PML siRNA (h): sc-36284, PML shRNA Plasmid (h): sc-36284-SH and PML shRNA (h) Lentiviral Particles: sc-36284-V.


Molecular Weight of PML isoforms: 78/97 kDa.

Positive Controls: HeLa whole cell lysate: sc-2200, K-562 nuclear extract: sc-2130 or K-562 whole cell lysate: sc-2203.


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850. 4) Immunohistochemistry: use m-IgG κ BP-HRP: sc-516102 with DAB, 50X: sc-24982 and Immunohistomount: sc-45086, or Organo/Limonene Mount: sc-45087.

DATA



PML (5G191): sc-71910. Western blot analysis of PML expression in HeLa whole cell lysate.



PML (5G191): sc-71910. Immunoperoxidase staining of formalin fixed, paraffin-embedded human lymph node tissue showing nuclear staining of cells in germinal center (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human urinary bladder tissue showing nuclear staining of urothelial cells (B).

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.



See **PML (E-11): sc-377390** for PML antibody conjugates, including AC, HRP, FITC, PE, and Alexa Fluor® 488, 546, 594, 647, 680 and 790.