

Rtn-1/2 (4A69): sc-71981

BACKGROUND

The reticulon (Rtn) family consists of four members: Rtn-1 (also designated neuroendocrine-specific protein or NSP), Rtn-2 (also designated NSP-like-1), Rtn-3 (also designated NSP-like-2) and Nogo (also designated Rtn-4A). Localized on human chromosome 14q23.1, the gene encoding Rtn-1 is expressed as three isoforms: Rtn-1A (NSP-A), Rtn-1B (NSP-B) and Rtn-1C (NSP-C), which are produced by alternative splicing, are anchored to the endoplasmic reticulum in neural and neuroendocrine tissues and cells, and may be involved in neuroendocrine secretion or in membrane trafficking. The gene encoding human Rtn-2 is located on chromosome 19q13.32 and also encodes three isoforms. Rtn-2-A and Rtn-2-C are produced by the use of alternative initiation sites, whereas Rtn-2-B is an alternative splice variant of the Rtn-2-A isoform. Rtn-2-A and Rtn-2-B are highly expressed in brain, while Rtn-2-C is primarily expressed in skeletal muscle. The human Rtn-3 gene is located on chromosome 11q13 and is widely expressed, with the highest expression being in brain.

REFERENCES

- Senden, N.H., et al. 1994. Subcellular localization and supramolecular organization of neuroendocrine-specific protein B (NSP-B) in small cell lung cancer. *Eur. J. Cell Biol.* 65: 341-353.
- van de Velde, H.J., et al. 1994. NSP-encoded reticulons are neuroendocrine markers of a novel category in human lung cancer diagnosis. *Cancer Res.* 54: 4769-4776.
- Geisler, J.G., et al. 1998. Molecular cloning of a novel mouse gene with predominant muscle and neural expression. *Mamm. Genome* 9: 274-282.
- Roebroek, A.J., et al. 1998. CDNA cloning, genomic organization, and expression of the human RTN2 gene, a member of a gene family encoding reticulons. *Genomics* 51: 98-106.
- Hens, J., et al. 1998. Neuronal differentiation is accompanied by NSP-C expression. *Cell Tissue Res.* 292: 229-237.
- Moreira, E.F., et al. 1999. Cloning of a novel member of the reticulon gene family (RTN3): gene structure and chromosomal localization to 11q13. *Genomics* 58: 73-81.
- GrandPré, T., et al. 2000. Identification of the Nogo inhibitor of axon regeneration as a reticulon protein. *Nature* 403: 439-444.

CHROMOSOMAL LOCATION

Genetic locus: RTN1 (human) mapping to 14q23.1, RTN2 (human) mapping to 19q13.32.

SOURCE

Rtn-1/2 (4A69) is a mouse monoclonal antibody raised against the small cell lung cancer cell line NCI-H82.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

APPLICATIONS

Rtn-1/2 (4A69) is recommended for detection of Rtn-1 and Rtn-2 of human, rabbit and Rhesus monkey origin by Western Blotting (starting dilution 1:100, dilution range 1:100-1:1,000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Molecular Weight of Rtn-1: 100 kDa.


Molecular Weight of Rtn-2: 58 kDa.

Positive Controls: Rtn-1 (h): 293T Lysate: sc-111610.

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

DATA



Rtn-1/2 (4A69): sc-71981. Western blot analysis of Rtn-1 expression in non-transfected: sc-117752 (A) and human Rtn-1 transfected: sc-111610 (B) 293T whole cell lysates.

SELECT PRODUCT CITATIONS

- Liang, C., et al. 2019. Single-cell sequencing of iPSC-dopamine neurons reconstructs disease progression and identifies HDAC4 as a regulator of Parkinson cell phenotypes. *Cell Stem Cell* 24: 93-106.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.