

MCP-3 siRNA (h): sc-72035

BACKGROUND

Eotaxin and the monocyte chemotactic proteins, MCP-1-5, form a subfamily of the C-C (or β) chemokines, which are characterized by a set of conserved adjacent cysteines. MCPs are produced by a variety of cells, including T lymphocytes, subsequent to their activation with cytokines such as IL-1, TNF α and IFN- γ . *In vitro* studies have shown that the MCP isoforms exhibit their chemotactic effects on different subpopulations of lymphocytes. MCP-3 has been shown to have the broadest range of influence, activating monocytes, dendritic cells, lymphocytes, NK cells, eosinophils, basophils and neutrophils.

REFERENCES

1. Charo, I.F., et al. 1994. Molecular cloning and functional expression of two monocyte chemoattractant protein 1 receptors reveals alternative splicing of the carboxyl-terminal tails. *Proc. Natl. Acad. Sci. USA* 91: 2752-2756.
2. Taub, D.D., et al. 1995. Monocyte chemotactic protein-1 (MCP-1), -2 and -3 are chemotactic for human T lymphocytes. *J. Clin. Invest.* 95: 1370-1376.
3. Weber, M., et al. 1995. Monocyte chemotactic protein MCP-2 activates human basophil and eosinophil leukocytes similar to MCP-3. *J. Immunol.* 154: 4166-4172.
4. Combadiere, C., et al. 1995. Monocyte chemoattractant protein-3 is a functional ligand for CC chemokine receptors 1 and 2B. *J. Biol. Chem.* 270: 29671-29675.
5. Proost, P., et al. 1996. Human monocyte chemotactic proteins-2 and -3: structural and functional comparison with MCP-1. *J. Leukoc. Biol.* 59: 67-74.
6. Dubois, P.M., et al. 1996. Early signal transduction by the receptor to the chemokine monocyte chemotactic protein-1 in a murine T cell hybrid. *J. Immunol.* 156: 1356-1361.
7. Beall, C.J., et al. 1996. Site-directed mutagenesis of monocyte chemoattractant protein-1 identifies two regions of the polypeptide essential for biological activity. *Biochem. J.* 313: 633-640.
8. Kuna, P., et al. 1996. Chemokines in seasonal allergic rhinitis. *J. Allergy Clin. Immunol.* 97: 104-112.

CHROMOSOMAL LOCATION

Genetic locus: CCL7 (human) mapping to 17q12.

PRODUCT

MCP-3 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see MCP-3 shRNA Plasmid (h): sc-72035-SH and MCP-3 shRNA (h) Lentiviral Particles: sc-72035-V as alternate gene silencing products.

For independent verification of MCP-3 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-72035A, sc-72035B and sc-72035C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

MCP-3 siRNA (h) is recommended for the inhibition of MCP-3 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

MCP-3 (G-2): sc-515890 is recommended as a control antibody for monitoring of MCP-3 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz MarkerTM Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor MCP-3 gene expression knockdown using RT-PCR Primer: MCP-3 (h)-PR: sc-72035-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.