

Ly-49F siRNA (m): sc-72051

BACKGROUND

The activity of natural killer (NK) cells is regulated by members of multiple receptor families that recognize class I MHC molecules, such as the killer cell inhibitory receptor/leukocyte immunoglobulin-like receptor (KIR/LIR) family and the C-type lectin superfamily. The KIR/LIR family includes p91A (also designated pp130 or PIR-B, for paired immunoglobulin-like receptor-B) and p91B (also designated PIR-A). p91A acts as an inhibitory receptor through interactions with SHP-1, whereas p91B acts as an activating receptor. CD94, NKG2 and Ly-49 are members of the C-type lectin superfamily of type II membrane glycoproteins. CD94 forms heterodimers with NKG2 isoforms on the surface of NK cells, whereas Ly-49 isoforms form homodimers. There are at least 21 Ly-49 proteins, ranging from 230 to 280 amino acids in length. The Ly-49 isoforms, which are clustered on human chromosome 16, all function to present the C-type lectin domain on the cell membrane. NKG2-D, expressed on NK cells, $\gamma\delta$ T cells, and CD8 $^{+}$ $\alpha\beta$ T cells, is a receptor for the stress inducible protein MICA, an antigen frequently expressed in epithelial tumors.

REFERENCES

1. Junggren, H.G., et al. 1990. In search of the "missing self": MHC molecules and NK cell recognition. *Immunol. Today* 11: 237-244.
2. Ortaldo, J.R., et al. 1998. The Ly-49 family: regulation of cytotoxicity and cytokine production in murine CD3 $^{+}$ cells. *J. Immunol.* 160: 1158-1165.
3. Robson MacDonald, H., et al. 1998. Developmentally regulated ext and selection of NK1.1 $^{+}$ T cells. *J. Exp. Med.* 187: 2109-2124.
4. Manilay, J.O., et al. 1999. Levels of Ly-49 receptor expression are determined by the frequency of interactions with MHC ligands: evidence against receptor calibration to a "useful" level. *J. Immunol.* 163: 2628-2633.
5. Naper, C., et al. 1999. Genes in two major histocompatibility complex class I regions control selection, phenotype, and function of a rat Ly-49 natural killer cell subset. *Eur. J. Immunol.* 29: 2046-2053.
6. Coles, M.C., et al. 2000. Memory CD8 T lymphocytes express inhibitory MHC-specific Ly49 receptors. *Eur. J. Immunol.* 30: 236-244.
7. Kane, K.P., et al. 2001. Specificity and function of activating Ly-49 receptors. *Immunol. Rev.* 181: 104-114.

CHROMOSOMAL LOCATION

Genetic locus: Klra6 (mouse) mapping to 6 F3.

PRODUCT

Ly-49F siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Ly-49F shRNA Plasmid (m): sc-72051-SH and Ly-49F shRNA (m) Lentiviral Particles: sc-72051-V as alternate gene silencing products.

For independent verification of Ly-49F (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-72051A, sc-72051B and sc-72051C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Ly-49F siRNA (m) is recommended for the inhibition of Ly-49F expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Ly-49F (HBF-719): sc-53023 is recommended as a control antibody for monitoring of Ly-49F gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Ly-49F gene expression knockdown using RT-PCR Primer: Ly-49F (m)-PR: sc-72051-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.