

# COL1A2 siRNA (h): sc-72156

## BACKGROUND

The extensive family of COL gene products (collagens) is composed of several chain types, including fibril-forming interstitial collagens (types I, II, III and V) and basement membrane collagens (type IV), each type containing multiple isoforms. Collagens are fibrous, extracellular matrix proteins with high tensile strength and are the major components of connective tissue, such as tendons and cartilage. All collagens contain a triple helix domain and frequently show lateral self-association in order to form complex connective tissues. Several collagens also play a role in cell adhesion, important for maintaining normal tissue architecture and function.

## REFERENCES

1. Bateman, J.F., et al. 1996. Collagen superfamily. In Comper, W.D., ed., *Extracellular Matrix, Vol. 2: Molecular Components and Interactions*. Amsterdam: Harwood Academic Publishers, 22-67.
2. McCarthy, J.B., et al. 1996. Cell adhesion to collagenous matrices. *Biopolymers* 40: 371-381.
3. Myers, L.K., et al. 1997. Collagen-induced arthritis, an animal model of autoimmunity. *Life Sci.* 61: 1861-1878.
4. Staines, N.A., et al. 1997. Nasal tolerance to dominant and subdominant epitopes of Collagen Type II and protection against collagen-induced arthritis. *Biochem. Soc. Trans.* 25: 661-664.
5. Engel, J. 1997. Versatile collagens in invertebrates. *Science* 277: 1785-1786.

## CHROMOSOMAL LOCATION

Genetic locus: COL1A2 (human) mapping to 7q21.3.

## PRODUCT

COL1A2 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see COL1A2 shRNA Plasmid (h): sc-72156-SH and COL1A2 shRNA (h) Lentiviral Particles: sc-72156-V as alternate gene silencing products.

For independent verification of COL1A2 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-72156A, sc-72156B and sc-72156C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

COL1A2 siRNA (h) is recommended for the inhibition of COL1A2 expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

Pro-COL1A2 (D-6): sc-166572 is recommended as a control antibody for monitoring of COL1A2 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\lambda$  BP-HRP: sc-516132 or m-IgG $\lambda$  BP-HRP (Cruz Marker): sc-516132-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\lambda$  BP-FITC: sc-516185 or m-IgG $\lambda$  BP-PE: sc-516186 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor COL1A2 gene expression knockdown using RT-PCR Primer: COL1A2 (h)-PR: sc-72156-PR (20  $\mu$ l, 583 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## SELECT PRODUCT CITATIONS

1. Nacu, N., et al. 2008. Macrophages produce TGF- $\beta$ -induced ( $\beta$ -ig-h3) following ingestion of apoptotic cells and regulate MMP14 levels and collagen turnover in fibroblasts. *J. Immunol.* 180: 5036-5044.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.