

CLEC-5A siRNA (m): sc-72177

BACKGROUND

The C-type lectin/C-type lectin-like domain (CTL/CTLD) superfamily consists of a variety of proteins that share a common protein fold and have diverse functions, including cell-cell signaling, cell adhesion, glycoprotein turnover and immune responses. CLEC-5A (C-type lectin domain family 5 member A), also known as MDL-1 (myeloid DAP12-associating lectin) or CLECSF5, is a 188 amino acid single-pass type II membrane protein that contains one C-type lectin domain and belongs to the C-type lectin superfamily. Expressed in peripheral blood monocytes, CLEC-5A is thought to be involved in the pro-inflammatory activation of myeloid cells, specifically by mediating DAP12 signaling in a calcium-dependent manner. The signaling cascade in which CLEC-5A is involved is associated with dengue hemorrhagic fever and dengue shock syndrome, suggesting a role for CLEC-5A in dengue viral host infection. Multiple isoforms of CLEC-5A exist due to alternative splicing events.

REFERENCES

- Drickamer, K. 1999. C-type lectin-like domains. *Curr. Opin. Struct. Biol.* 9: 585-590.
- Bakker, A.B., et al. 1999. Myeloid DAP12-associating lectin (MDL)-1 is a cell surface receptor involved in the activation of myeloid cells. *Proc. Natl. Acad. Sci. USA* 96: 9792-9796.
- Yim, D., et al. 2001. Molecular cloning and expression pattern of porcine myeloid DAP12-associating lectin-1. *Cell. Immunol.* 209: 42-48.
- Gingras, et al. 2002. TREM-1, MDL-1, and DAP12 expression is associated with a mature stage of myeloid development. *Mol. Immunol.* 38: 817-824.
- Online Mendelian Inheritance in Man, OMIM[™]. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 604987. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Ebner, S., et al. 2003. Evolutionary analysis reveals collective properties and specificity in the C-type lectin and lectin-like domain superfamily. *Proteins* 53: 44-55.
- Aoki, N., et al. 2004. Differential regulation of DAP12 and molecules associated with DAP12 during host responses to mycobacterial infection. *Infect. Immun.* 72: 2477-2483.

CHROMOSOMAL LOCATION

Genetic locus: Clec5a (mouse) mapping to 6 B1.

PRODUCT

CLEC-5A siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see CLEC-5A shRNA Plasmid (m): sc-72177-SH and CLEC-5A shRNA (m) Lentiviral Particles: sc-72177-V as alternate gene silencing products.

For independent verification of CLEC-5A (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-72177A, sc-72177B and sc-72177C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

CLEC-5A siRNA (m) is recommended for the inhibition of CLEC-5A expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CLEC-5A gene expression knockdown using RT-PCR Primer: CLEC-5A (m)-PR: sc-72177-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.