

ADAM29 siRNA (h): sc-72237

BACKGROUND

The ADAM (a disintegrin and metalloprotease) protein family, which includes over 30 membrane-anchored, glycosylated, Zn^{2+} dependent proteases, plays a role in cell-cell and cell-matrix interface related processes, including fertilization, muscle fusion, secretion of tumor necrosis factor- α (TNF α), and modulation of the neurogenic function of Notch and Delta. The ADAM proteins possess a signal-domain, a pro-domain, a metalloprotease domain, a disintegrin domain (Integrin ligand), a cysteine-rich region, an epidermal growth factor-like domain, a transmembrane domain, and a cytoplasmic tail. ADAMs are expressed in a wide range of mammalian tissues and several are abundantly expressed in the male reproductive tract. Both ADAM29 and ADAM30 show testis-specific expression. The genes encoding human ADAM29 and ADAM30 map to chromosomes 4q34.1 and 1p12, respectively, and encode multiple isoforms of each protein.

REFERENCES

1. Wolfsberg, T.G., et al. 1995. ADAM, a novel family of membrane proteins containing a disintegrin and metallo-protease domain: multipotential functions in cell-cell and cell-matrix interactions. *J. Cell Biol.* 131: 275-278.
2. Stone, A.L., et al. 1999. Structure-function analysis of the ADAM family of disintegrin-like and metalloproteinase-containing proteins. *J. Protein Chem.* 18: 447-465.
3. Xu, R., et al. 1999. Molecular cloning and mapping of a novel ADAM gene (ADAM29) to human chromosome 4. *Genomics* 62: 537-539.
4. Cerretti, D.P., et al. 1999. Isolation of two novel metalloproteinase-disintegrin (ADAM) cDNAs that show testis-specific gene expression. *Biochem. Biophys. Res. Commun.* 263: 810-815.
5. Primakoff, P. and Myles, D.G. 2000. The ADAM gene family: surface proteins with adhesion and protease activity. *Trends Genet.* 16: 83-87.

CHROMOSOMAL LOCATION

Genetic locus: ADAM29 (human) mapping to 4q34.1.

PRODUCT

ADAM29 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ADAM29 shRNA Plasmid (h): sc-72237-SH and ADAM29 shRNA (h) Lentiviral Particles: sc-72237-V as alternate gene silencing products.

For independent verification of ADAM29 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-72237A, sc-72237B and sc-72237C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ADAM29 siRNA (h) is recommended for the inhibition of ADAM29 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ADAM29 gene expression knockdown using RT-PCR Primer: ADAM29 (h)-PR: sc-72237-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.