

Liprin α 1 siRNA (m): sc-72331

BACKGROUND

Liprins interact with members of the leukocyte common antigen-related (LAR) family of transmembrane protein tyrosine phosphatases, which are implicated in axon guidance and mammary gland development. Liprins are multivalent proteins that form complex structures and act as scaffolds for the recruitment and anchoring of LAR phosphatases. Based on sequence similarities and binding characteristics, liprins are subdivided into α and β liprins. Both α and β liprins homodimerize via their N-terminal, coiled-coil regions. Liprin α 1 is a ubiquitously expressed protein that interacts with the tumor suppressor ING4 to regulate cell migration and possibly prevent metastasis. The interaction between LAR and Liprin α 1 can be weakened by treatment of Liprin α 1 with calf intestinal phosphatase.

REFERENCES

1. Serra-Pagès, C., et al. 1998. Liprins, a family of LAR transmembrane protein-tyrosine phosphatase-interacting proteins. *J. Biol. Chem.* 273: 15611-15620.
2. Kaufmann, N., et al. 2002. *Drosophila* Liprin- α and the receptor phosphatase Dlar control synapse morphogenesis. *Neuron* 34: 27-38.
3. Ko, J., et al. 2003. Interaction between Liprin- α and GIT1 is required for AMPA receptor targeting. *J. Neurosci.* 23: 1667-1677.
4. Patel, M.R., et al. 2006. Hierarchical assembly of presynaptic components in defined *C. elegans* synapses. *Nat. Neurosci.* 9: 1488-1498.
5. Unoki, M., et al. 2006. Novel splice variants of ING4 and their possible roles in the regulation of cell growth and motility. *J. Biol. Chem.* 281: 34677-34686.
6. Hofmeyer, K., et al. 2006. Liprin- α has LAR-independent functions in R7 photoreceptor axon targeting. *Proc. Natl. Acad. Sci. USA* 103: 11595-11600.
7. Choe, K.M., et al. 2006. Liprin- α is required for photoreceptor target selection in *Drosophila*. *Proc. Natl. Acad. Sci. USA* 103: 11601-11606.
8. Olsen, O., et al. 2006. Synaptic transmission regulated by a presynaptic MALS/Liprin- α protein complex. *Curr. Opin. Cell Biol.* 18: 223-227.

CHROMOSOMAL LOCATION

Genetic locus: Ppfia1 (mouse) mapping to 7 F5.

PRODUCT

Liprin α 1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Liprin α 1 shRNA Plasmid (m): sc-72331-SH and Liprin α 1 shRNA (m) Lentiviral Particles: sc-72331-V as alternate gene silencing products.

For independent verification of Liprin α 1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-72331A, sc-72331B and sc-72331C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Liprin α 1 siRNA (m) is recommended for the inhibition of Liprin α 1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Liprin α 1 (A-6): sc-398030 is recommended as a control antibody for monitoring of Liprin α 1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Liprin α 1 gene expression knockdown using RT-PCR Primer: Liprin α 1 (m)-PR: sc-72331-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.