

PIG-B siRNA (h): sc-72355

BACKGROUND

Phosphatidylinositol-glycans (PIGs) are multi-pass transmembrane proteins that localize to the endoplasmic reticulum. PIGs exhibit various functions but all are crucial for the biosynthesis of the glycosylphosphatidylinositol (GPI)-anchor. Some PIG proteins are components of the GPI transamidase complex and play a role in the recognition of either the GPI attachment signal or the lipid portion of GPI. Other PIGs belong to the glycosyltransferase complex and function in the transfer of N-acetylglucosamine (GlcNAc) to phosphatidylinositol (PI). A variety of other PIGs play distinct roles in GPI synthesis including mannosylation of the GPI-anchor. PIG-B, an α 1,2-mannosyltransferase, is also referred to as GPI mannosyltransferase III (GPI-MT-III). It is responsible for adding the third mannose in the synthesis of the GPI anchor. PIG-B transfers the mannose from dolichol-phosphate-mannose (Dol-P-Man) and for this reason, it is a member of the Dol-P-Man-dependent mannosyltransferase family.

REFERENCES

1. Takahashi, M., Inoue, N., Ohishi, K., Maeda, Y., Nakamura, N., Endo, Y., Fujita, T., Takeda, J. and Kinoshita, T. 1996. PIG-B, a membrane protein of the endoplasmic reticulum with a large luminal domain, is involved in transferring the third mannose of the GPI anchor. *EMBO J.* 15: 4254-4261.
2. Nagamune, K., Nozaki, T., Maeda, Y., Ohishi, K., Fukuma, T., Hara, T., Schwarz, R.T., Sutterlin, C., Brun, R., Riezman, H. and Kinoshita, T. 2000. Critical roles of glycosylphosphatidylinositol for *Trypanosoma brucei*. *Proc. Natl. Acad. Sci. USA* 97: 10336-10341.
3. Maeda, Y., Watanabe, R., Harris, C.L., Hong, Y., Ohishi, K., Kinoshita, K. and Kinoshita, T. 2001. PIG-M transfers the first mannose to glycosylphosphatidylinositol on the luminal side of the ER. *EMBO J.* 20: 250-261.
4. Grimme, S.J., Westfall, B.A., Wiedman, J.M., Taron, C.H. and Orlean, P. 2001. The essential Smp3 protein is required for addition of the side-branching fourth mannose during assembly of yeast glycosylphosphatidylinositols. *J. Biol. Chem.* 276: 27731-27739.
5. Delorenzi, M., Sexton, A., Shams-Eldin, H., Schwarz, R.T., Speed, T. and Schofield, L. 2002. Genes for glycosylphosphatidylinositol toxin biosynthesis in *Plasmodium falciparum*. *Infect. Immun.* 70: 4510-4522.
6. Oriol, R., Martinez-Duncker, I., Chantret, I., Mollicone, R. and Codogno, P. 2002. Common origin and evolution of glycosyltransferases using Dol-P-monosaccharides as donor substrate. *Mol. Biol. Evol.* 19: 1451-1463.
7. Ikezawa, H. 2002. Glycosylphosphatidylinositol (GPI)-anchored proteins. *Biol. Pharm. Bull.* 25: 409-417.
8. Kang, J.Y., Hong, Y., Ashida, H., Shishioh, N., Murakami, Y., Morita, Y.S., Maeda, Y. and Kinoshita, T. 2005. PIG-V involved in transferring the second mannose in glycosylphosphatidylinositol. *J. Biol. Chem.* 280: 9489-9497.
9. Wiedman, J.M., Fabre, A.L., Taron, B.W., Taron, C.H. and Orlean, P. 2007. *In vivo* characterization of the GPI assembly defect in yeast *mcd4-174* mutants and bypass of the *Mcd4p*-dependent step in *mcd4Delta* cells. *FEMS Yeast Res.* 7: 78-83.

CHROMOSOMAL LOCATION

Genetic locus: PIGB (human) mapping to 15q21.3.

PRODUCT

PIG-B siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PIG-B shRNA Plasmid (h): sc-72355-SH and PIG-B shRNA (h) Lentiviral Particles: sc-72355-V as alternate gene silencing products.

For independent verification of PIG-B (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-72355A, sc-72355B and sc-72355C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PIG-B siRNA (h) is recommended for the inhibition of PIG-B expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PIG-B gene expression knockdown using RT-PCR Primer: PIG-B (h)-PR: sc-72355-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.