

PIG-C siRNA (m): sc-72358

BACKGROUND

Phosphatidylinositol-glycans (PIGs) are multi-pass transmembrane proteins that localize to the endoplasmic reticulum. PIGs exhibit various functions but all are crucial for the biosynthesis of the glycosylphosphatidylinositol (GPI)-anchor. Some PIG proteins are components of the GPI transamidase complex and play a role in the recognition of either the GPI attachment signal or the lipid portion of GPI. Other PIGs belong to the glycosyltransferase complex (GPI-N-acetylglucosaminyltransferase or GPI-GnT) and function in the transfer of N-acetylglucosamine (GlcNAc) to phosphatidylinositol (PI). A variety of other PIGs play distinct roles in GPI synthesis. PIG-C is a component of the GPI-GnT complex which is responsible for the first step in GPI synthesis, the transfer of GlcNAc to PI from UDP-GlcNAc. PIG-C may function as a scaffold protein, anchoring the complex to the ER and facilitating the interaction among other subunits. A mutation in the gene encoding PIG-C may result in the absence of GPI-anchored proteins.

REFERENCES

1. Inoue, N., et al. 1996. PIG-C, one of the three human genes involved in the first step of glycosylphosphatidylinositol biosynthesis is a homologue of *Saccharomyces cerevisiae* GPI2. *Biochem. Biophys. Res. Commun.* 226: 193-199.
2. Watanabe, R., et al. 1998. The first step of glycosylphosphatidylinositol biosynthesis is mediated by a complex of PIG-A, PIG-H, PIG-C and GPI1. *EMBO J.* 17: 877-885.
3. Watanabe, R., et al. 2000. Initial enzyme for glycosylphosphatidylinositol biosynthesis requires PIG-P and is regulated by DPM2. *EMBO J.* 19: 4402-4411.
4. Tiede, A., et al. 2000. Characterisation of the enzymatic complex for the first step in glycosylphosphatidylinositol biosynthesis. *Int. J. Biochem. Cell Biol.* 32: 339-350.
5. Ikezawa, H. 2002. Glycosylphosphatidylinositol (GPI)-anchored proteins. *Biol. Pharm. Bull.* 25: 409-417.
6. Newman, H.A., et al. 2005. Gpi19, the *Saccharomyces cerevisiae* homologue of mammalian PIG-P, is a subunit of the initial enzyme for glycosylphosphatidylinositol anchor biosynthesis. *Eukaryotic Cell* 4: 1801-1807.
7. Maneesri, J., et al. 2005. Deletion of MCD 4 involved in glycosylphosphatidylinositol (GPI) anchor synthesis leads to an increase in β -1,6-glucan level and a decrease in GPI-anchored protein and mannan levels in the cell wall of *Saccharomyces cerevisiae*. *J. Biosci. Bioeng.* 99: 354-360.
8. Li, H., et al. 2007. Glycosylphosphatidylinositol (GPI) anchor is required in *Aspergillus fumigatus* for morphogenesis and virulence. *Mol. Microbiol.* 64: 1014-1027.
9. Pittet, M. and Conzelmann, A. 2007. Biosynthesis and function of GPI proteins in the yeast *Saccharomyces cerevisiae*. *Biochim. Biophys. Acta* 1771: 405-420.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: *Pigc* (mouse) mapping to 1 H2.1.

PRODUCT

PIG-C siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PIG-C shRNA Plasmid (m): sc-72358-SH and PIG-C shRNA (m) Lentiviral Particles: sc-72358-V as alternate gene silencing products.

For independent verification of PIG-C (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-72358A, sc-72358B and sc-72358C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PIG-C siRNA (m) is recommended for the inhibition of PIG-C expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PIG-C gene expression knockdown using RT-PCR Primer: PIG-C (m)-PR: sc-72358-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.