

# ADAMTS-17 siRNA (m): sc-72446

## BACKGROUND

ADAMTS (a disintegrin and metalloproteinase domain with Thrombospondin 1 modules) is a family of zinc-dependent proteases that are implicated in a variety of normal and pathological conditions, including arthritis and cancer. ADAMTS protein family members contain an N-terminal propeptide domain, a metalloproteinase domain, a disintegrin-like domain and a C-terminus that contains a varying number of Thrombospondin 1 (TSP-1) motifs. ADAMTS-17 (ADAM metalloproteinase with Thrombospondin 1 motif, 17) is a 1,095 amino acid secreted protein that belongs to the ADAMTS family. Expressed in liver, prostate, brain and in fetal lung, ADAMTS-17 has seven N-glycosylation sites and shares 56% sequence identity with a related family member, ADAMTS-19. ADAMTS-17 has a conserved cysteine residue in its cysteine-switch motif that, when bound to one zinc ion, inhibits the protein's enzymatic activity.

## REFERENCES

1. Kuno, K., et al. 1997. Molecular cloning of a gene encoding a new type of metalloproteinase-disintegrin family protein with Thrombospondin motifs as an inflammation associated gene. *J. Biol. Chem.* 272: 556-562.
2. Tang, B.L. and Hong, W. 1999. ADAMTS: a novel family of proteases with an ADAM protease domain and Thrombospondin 1 repeats. *FEBS Lett.* 445: 223-225.
3. Cal, S., et al. 2002. Cloning, expression analysis, and structural characterization of seven novel human ADAMTSs, a family of metalloproteinases with disintegrin and Thrombospondin 1 domains. *Gene* 283: 49-62.
4. Online Mendelian Inheritance in Man, OMIM<sup>™</sup>. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 607511. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Overall, C.M., et al. 2004. Protease degradomics: mass spectrometry discovery of protease substrates and the CLIP-CHIP, a dedicated DNA microarray of all human proteases and inhibitors. *Biol. Chem.* 385: 493-504.
6. Davidson, R.K., et al. 2006. Expression profiling of metalloproteinases and their inhibitors in synovium and cartilage. *Arthritis Res. Ther.* 8: R124.

## CHROMOSOMAL LOCATION

Genetic locus: Adamts17 (mouse) mapping to 7 C.

## PRODUCT

ADAMTS-17 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ADAMTS-17 shRNA Plasmid (m): sc-72446-SH and ADAMTS-17 shRNA (m) Lentiviral Particles: sc-72446-V as alternate gene silencing products.

For independent verification of ADAMTS-17 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-72446A, sc-72446B and sc-72446C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

ADAMTS-17 siRNA (m) is recommended for the inhibition of ADAMTS-17 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

ADAMTS-17 (Q-12): sc-100480 is recommended as a control antibody for monitoring of ADAMTS-17 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ADAMTS-17 gene expression knockdown using RT-PCR Primer: ADAMTS-17 (m)-PR: sc-72446-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.