



Angiomotin-L1 siRNA (m): sc-72492

BACKGROUND

Angiomotin, also known as AMOT, is a 1,084 amino acid protein that belongs to the motin family of angiostatin binding proteins. Members of the motin family contain conserved coiled-coil domains and PDZ binding motifs at their C-termini. Expressed in skeletal muscle and placenta, Angiomotin localizes to the cell surface at tight junctions and is believed to be involved in tight junction maintenance. Angiomotin binds to angiostatin and plays a vital role in angiogenesis, promoting tubule formation and growth factor-induced migration of endothelial cells. This suggests that Angiomotin may be an important player in tumor angiogenesis and could serve as a potential therapeutic target in cancer. Angiomotin-L1 (Angiomotin like 1), also known as AMOTL1 or JEAP, is a 956 amino acid peripheral membrane protein that localizes to tight junctions and may help to control cellular permeability and to maintain cell polarity.

REFERENCES

1. Kikuno, R., et al. 1999. Prediction of the coding sequences of unidentified human genes. XIV. The complete sequences of 100 new cDNA clones from brain which code for large proteins *in vitro*. DNA Res. 6: 197-205.
2. Troyanovsky, B., et al. 2001. Angiomotin: an angiostatin binding protein that regulates endothelial cell migration and tube formation. J. Cell Biol. 152: 1247-1254.
3. Bratt, A., et al. 2002. Angiomotin belongs to a novel protein family with conserved coiled-coil and PDZ binding domains. Gene 298: 69-77.
4. Wells, C.D., et al. 2006. A Rich1/Amot complex regulates the Cdc42 GTPase and apical-polarity proteins in epithelial cells. Cell 125: 535-548.
5. Ernkvist, M., et al. 2006. p130-Angiomotin associates to Actin and controls endothelial cell shape. FEBS J. 273: 2000-2011.
6. Huang, H., et al. 2007. Amotl2 is essential for cell movements in zebrafish embryo and regulates c-Src translocation. Development 134: 979-988.
7. Aase, K., et al. 2007. Angiomotin regulates endothelial cell migration during embryonic angiogenesis. Genes Dev. 21: 2055-2068.

CHROMOSOMAL LOCATION

Genetic locus: Amotl1 (mouse) mapping to 9 A1.

PRODUCT

Angiomotin-L1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Angiomotin-L1 shRNA Plasmid (m): sc-72492-SH and Angiomotin-L1 shRNA (m) Lentiviral Particles: sc-72492-V as alternate gene silencing products.

For independent verification of Angiomotin-L1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-72492A, sc-72492B and sc-72492C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Angiomotin-L1 siRNA (m) is recommended for the inhibition of Angiomotin-L1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Angiomotin-L1 gene expression knockdown using RT-PCR Primer: Angiomotin-L1 (m)-PR: sc-72492-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.