SANTA CRUZ BIOTECHNOLOGY, INC.

Chibby siRNA (h): sc-72890



BACKGROUND

Chibby, also known as cytosolic leucine-rich protein or PIGEA-14, is a 126 amino acid highly conserved protein that inhibits β -catenin-mediated transcriptional activation by competing with LEF-1 to bind β -catenin. Chibby may also play a role in the regulation of the intracellular location of Polycystin-2 and other intracellular proteins. Acting as a homodimer, Chibby is subcellularly localized to the nucleus and golgi apparatus within the *trans*-Golgi network. Interaction with 14-3-3 results in the sequestration of Chibby to the cytoplasm and the formation of a stable complex with β -catenin, thereby facilitating nuclear export of β -catenin. Though widely expressed, Chibby is found at highest levels in skeletal muscle, heart, placenta and kidney. Down-regulation of Chibby is observed in thyroid and metastatic uterine tumors, suggesting that the gene encoding Chibby may function as a tumor suppressor.

REFERENCES

- 1. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 607757. World Wide Web URL: http://www.ncbi.nlm.nih.gov/omim/
- 2. Takemaru, K., et al. 2003. Chibby, a nuclear β -catenin-associated antagonist of the Wnt/Wingless pathway. Nature 422: 905-909.
- 3. Gad, S., et al. 2004. Is the gene encoding Chibby implicated as a tumour suppressor in colorectal cancer ? BMC Cancer 4: 31.
- Hidaka, S., et al. 2004. PIGEA-14, a novel coiled-coil protein affecting the intracellular distribution of polycystin-2. J. Biol. Chem. 279: 35009-35016.
- 5. Jung, Y., et al. 2006. TC1 (C8orf4) enhances the Wnt/β-catenin pathway by relieving antagonistic activity of Chibby. Cancer Res. 66: 723-728.

CHROMOSOMAL LOCATION

Genetic locus: CBY1 (human) mapping to 22q13.1.

PRODUCT

Chibby siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Chibby shRNA Plasmid (h): sc-72890-SH and Chibby shRNA (h) Lentiviral Particles: sc-72890-V as alternate gene silencing products.

For independent verification of Chibby (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-72890A, sc-72890B and sc-72890C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNAse-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNAse-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Chibby siRNA (h) is recommended for the inhibition of Chibby expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Chibby (C-10): sc-393295 is recommended as a control antibody for monitoring of Chibby gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgGκ BP-FITC: sc-516140 or m-IgGκ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Chibby gene expression knockdown using RT-PCR Primer: Chibby (h)-PR: sc-72890-PR (20 μ I). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

 Singh, A. and Sen, E. 2017. Reciprocal role of SIRT6 and Hexokinase 2 in the regulation of autophagy driven monocyte differentiation. Exp. Cell Res. 360: 365-374.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.