



Xin β siRNA (m): sc-72934

BACKGROUND

Xin β , also known as CMYA3 (cardiomyopathy-associated protein 3) or XIRP2 (xin actin-binding repeat containing 2), is a 3,374 amino acid protein that co-localizes with Actin stress fibers at the cell junction and contains 28 Xin repeats. Interacting with F-Actin and α -actinin-2, Xin β functions to protect Actin filaments from depolymerization, thereby playing an important role in Actin organization and structural maintenance. Multiple isoforms of Xin β exist due to alternative splicing events. The gene encoding Xin β maps to human chromosome 2, which houses over 1,400 genes and comprises nearly 8% of the human genome. Harlequin ichthyosis, a rare and morbid skin deformity, is associated with mutations in the ABCA12 gene, while the lipid metabolic disorder sitosterolemia is associated with defects in the ABCG5 and ABCG8 genes. Additionally, an extremely rare recessive genetic disorder, Alström syndrome, is caused by mutations in the ALMS1 gene, which maps to chromosome 2.

REFERENCES

1. Sinn, H.W., Balsamo, J., Lilien, J. and Lin, J.J. 2002. Localization of the novel Xin protein to the adherens junction complex in cardiac and skeletal muscle during development. *Dev. Dyn.* 225: 1-13.
2. Pan, P.W., Li, K., Tuggle, C.K., Yu, M., Liu, B. and Zhao, S.H. 2003. Sequencing, tissue distribution and physical mapping of the porcine homologue of cardiomyopathy associated 3 (CMYA3). *Anim. Genet.* 34: 473-474.
3. Pacholsky, D., Vakeel, P., Himmel, M., Löwe, T., Stradal, T., Rottner, K., Fürst, D.O. and van der Ven, P.F. 2004. Xin repeats define a novel Actin-binding motif. *J. Cell Sci.* 117: 5257-5268.
4. Blasi, F., Bacchelli, E., Carone, S., Toma, C., Monaco, A.P., Bailey, A.J. and Maestrini, E. 2006. SLC25A12 and CMYA3 gene variants are not associated with autism in the IMGSAC multiplex family sample. *Eur. J. Hum. Genet.* 14: 123-126.
5. Huang, H.T., Brand, O.M., Mathew, M., Ignatiou, C., Ewen, E.P., McCalmon, S.A. and Naya, F.J. 2006. Myomaxin is a novel transcriptional target of MEF2A that encodes a Xin-related α -actinin-interacting protein. *J. Biol. Chem.* 281: 39370-39379.
6. Online Mendelian Inheritance in Man, OMIM™. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 609778. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: Xirp2 (mouse) mapping to 2 C1.3.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

Xin β siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Xin β shRNA Plasmid (m): sc-72934-SH and Xin β shRNA (m) Lentiviral Particles: sc-72934-V as alternate gene silencing products.

For independent verification of Xin β (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-72934A, sc-72934B and sc-72934C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Xin β siRNA (m) is recommended for the inhibition of Xin β expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Xin β gene expression knockdown using RT-PCR Primer: Xin β (m)-PR: sc-72934-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.