

COQ3 siRNA (m): sc-72974

BACKGROUND

Coenzyme Q (COQ), also referred to as ubiquinone, is a fat-soluble component of the electron transport chain that participates in aerobic cellular respiration within mitochondria and is essential for ATP-dependent energy production. COQ consists of a hydrophobic isoprenoid tail, which anchors it to the membrane, and a quinone head group, which is responsible for the activity of COQ in the respiratory chain. COQ3 (coenzyme Q3 homolog), also known as hexaprenyldihydroxybenzoate methyltransferase or DHHB methyltransferase (DHHB-MT), mitochondrial is a 369 amino acid protein that localizes to the mitochondrial matrix. COQ3 is a methyltransferase required for two steps in the biosynthesis of coenzyme Q.

REFERENCES

- Jonassen, T., et al. 2000. Isolation and functional expression of human COQ3, a gene encoding a methyltransferase required for ubiquinone biosynthesis. *J. Biol. Chem.* 275: 12381-12387.
- Wiemann, S., et al. 2001. Toward a catalog of human genes and proteins: sequencing and analysis of 500 novel complete protein coding human cDNAs. *Genome Res.* 11: 422-435.
- Online Mendelian Inheritance in Man, OMIM[™]. 2006. Johns Hopkins University, Baltimore, MD. MIM Number: 605196. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Hsieh, E.J., et al. 2007. *Saccharomyces cerevisiae* Coq9 polypeptide is a subunit of the mitochondrial coenzyme Q biosynthetic complex. *Arch. Biochem. Biophys.* 463: 19-26.
- Franke, B., et al. 2009. An association study of 45 folate-related genes in spina bifida: involvement of cubilin (CUBN) and tRNA aspartic acid methyltransferase 1 (TRDMT1). *Birth Defects Res. Part A Clin. Mol. Teratol.* 85: 216-226.

CHROMOSOMAL LOCATION

Genetic locus: Coq3 (mouse) mapping to 4 A3.

PRODUCT

COQ3 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see COQ3 shRNA Plasmid (m): sc-72974-SH and COQ3 shRNA (m) Lentiviral Particles: sc-72974-V as alternate gene silencing products.

For independent verification of COQ3 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-72974A, sc-72974B and sc-72974C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

COQ3 siRNA (m) is recommended for the inhibition of COQ3 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

COQ3 (E-2): sc-377469 is recommended as a control antibody for monitoring of COQ3 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor COQ3 gene expression knockdown using RT-PCR Primer: COQ3 (m)-PR: sc-72974-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.