

IFN- α / β R α (H-11): sc-7391

BACKGROUND

The type I interferons (IFNs), α and β , are a group of structurally and functionally related proteins that are induced by either viruses or double stranded RNA and defined by their ability to confer an antiviral state in cells. The α and β IFNs appear to compete with one another for binding to a common cell surface receptor, while immune IFN (IFN- γ) binds to a distinct receptor. The latter protein, IFN- α R, is only weakly responsive to type I interferons in contrast to IFN- α / β R, which binds to and responds effectively to IFN- β and to several of the IFN- α subtypes. Moreover, IFN- α / β R is physically associated with the cytoplasmic tyrosine kinase JAK1 and thus, in addition to ligand binding, appears to be functionally involved in signal transduction. The IFN- γ receptor complex consists of an α subunit (IFN- γ R α) and a β subunit that is 332 amino acids in length (mouse) and 337 amino acids in length (human).

CHROMOSOMAL LOCATION

Genetic locus: IFNAR1 (human) mapping to 21q22.11; Ifnar1 (mouse) mapping to 16 C3.3.

SOURCE

IFN- α / β R α (H-11) is a mouse monoclonal antibody raised against amino acids 458-557 mapping at the C-terminus (complete intracellular domain) of the IFN- α / β R α chain precursor of human origin.

PRODUCT

Each vial contains 200 μ g IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

IFN- α / β R α (H-11) is available conjugated to agarose (sc-7391 AC), 500 μ g/0.25 ml agarose in 1 ml, for IP; to HRP (sc-7391 HRP), 200 μ g/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-7391 PE), fluorescein (sc-7391 FITC), Alexa Fluor[®] 488 (sc-7391 AF488), Alexa Fluor[®] 546 (sc-7391 AF546), Alexa Fluor[®] 594 (sc-7391 AF594) or Alexa Fluor[®] 647 (sc-7391 AF647), 200 μ g/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor[®] 680 (sc-7391 AF680) or Alexa Fluor[®] 790 (sc-7391 AF790), 200 μ g/ml, for Near-Infrared (NIR) WB, IF and FCM.

APPLICATIONS

IFN- α / β R α (H-11) is recommended for detection of IFN- α / β R α chain of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)] and immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

Suitable for use as control antibody for IFN- α / β R α siRNA (h): sc-35637, IFN- α / β R α siRNA (m): sc-40090, IFN- α / β R α shRNA Plasmid (h): sc-35637-SH, IFN- α / β R α shRNA Plasmid (m): sc-40090-SH, IFN- α / β R α shRNA (h) Lentiviral Particles: sc-35637-V and IFN- α / β R α shRNA (m) Lentiviral Particles: sc-40090-V.

Molecular Weight of IFN- α subunit: 110 kDa.


Molecular Weight of IFN- β subunit: 95-100 kDa.

Molecular Weight of IFN- β subunit short form: 55 kDa.


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA



IFN- α / β R α (H-11): sc-7391. Western blot analysis of IFN- α / β R α expression in non-transfected: sc-117752 (A) and human IFN- α / β R α transfected: sc-113922 (B) 293T whole cell lysates.



IFN- α / β R α (H-11): sc-7391. Immunofluorescence staining of methanol-fixed HeLa cells showing membrane localization.

SELECT PRODUCT CITATIONS

- Miller, D.M., et al. 1999. Human cytomegalovirus inhibits IFN- α -stimulated antiviral and immunoregulatory responses by blocking multiple levels of IFN- α signal transduction. *J. Immunol.* 162: 6107-6113.
- Duan, X., et al. 2011. Differential roles for the interferon-inducible IFI16 and AIM2 innate immune sensors for cytosolic DNA in cellular senescence of human fibroblasts. *Mol. Cancer Res.* 9: 589-602.
- Lu, J., et al. 2012. Enterovirus 71 disrupts interferon signaling by reducing the level of interferon receptor 1. *J. Virol.* 86: 3767-3776.
- Heo, D.K., et al. 2012. Opposite regulatory effects of TRPC1 and TRPC5 on neurite outgrowth in PC12 cells. *Cell. Signal.* 24: 899-906.
- Liu, Y., et al. 2014. Enterovirus 71 inhibits cellular type I interferon signaling by downregulating JAK1 protein expression. *Viral Immunol.* 27: 267-276.
- Mathieu, M.G., et al. 2014. The helicase HAGE prevents interferon- α -induced PML expression in ABCB5⁺ malignant melanoma-initiating cells by promoting the expression of SOCS1. *Cell Death Dis.* 5: e1061.
- Han, T., et al. 2015. Set7 facilitates hepatitis C virus replication via enzymatic activity-dependent attenuation of the IFN-related pathway. *J. Immunol.* 194: 2757-2768.
- Zhang, M., et al. 2015. HSV-2 immediate-early protein US1 inhibits IFN- β production by suppressing association of IRF-3 with IFN- β promoter. *J. Immunol.* 194: 3102-3115.
- Bai, L., et al. 2015. Hepatitis B virus hijacks CTHRC1 to evade host immunity and maintain replication. *J. Mol. Cell Biol.* 7: 543-556.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

Alexa Fluor[®] is a trademark of Molecular Probes, Inc., Oregon, USA