

GPR101 siRNA (h): sc-75166

BACKGROUND

G protein-coupled receptors (GPRs), also known as seven transmembrane receptors, heptahelical receptors or 7TM receptors, comprise a superfamily of proteins that play a role in many different stimulus-response pathways. G protein-coupled receptors translate extracellular signals into intracellular signals (G protein activation) and they respond to a variety of signaling molecules, such as hormones and neurotransmitters. GPR101 (G protein-coupled receptor 101), also known as GPCR6, is a 508 amino acid multi-pass membrane protein that belongs to the G protein-coupled receptor 1 family. GPR101 functions as an orphan receptor that is thought to play a role in signaling events throughout the cell. Expressed predominantly in brain, GPR101 participates in a wide range of activities in the CNS via modulation of cAMP levels.

REFERENCES

1. Larhammar, D., Blomqvist, A.G. and Wahlestedt, C. 1993. The receptor revolution—multiplicity of G protein-coupled receptors. *Drug Des. Discov.* 9: 179-188.
2. Ji, T.H., Grossmann, M. and Ji, I. 1998. G protein-coupled receptors. I. Diversity of receptor-ligand interactions. *J. Biol. Chem.* 273: 17299-17302.
3. Lee, D.K., Nguyen, T., Lynch, K.R., Cheng, R., Vanti, W.B., Arkhitko, O., Lewis, T., Evans, J.F., George, S.R. and O'Dowd, B.F. 2001. Discovery and mapping of ten novel G protein-coupled receptor genes. *Gene* 275: 83-91.
4. Wittenberger, T., Schaller, H.C. and Hellebrand, S. 2001. An expressed sequence tag (EST) data mining strategy succeeding in the discovery of new G protein-coupled receptors. *J. Mol. Biol.* 307: 799-813.
5. Takeda, S., Kadowaki, S., Haga, T., Takaesu, H. and Mitaku, S. 2002. Identification of G protein-coupled receptor genes from the human genome sequence. *FEBS Lett.* 520: 97-101.
6. Bates, B., Zhang, L., Nawoschik, S., Kodangattil, S., Tseng, E., Kopsco, D., Kramer, A., Shan, Q., Taylor, N., Johnson, J., Sun, Y., Chen, H.M., Blatcher, M., Paulsen, J.E. and Pausch, M.H. 2006. Characterization of GPR101 expression and G protein coupling selectivity. *Brain Res.* 1087: 1-14.
7. Amisten, S., Braun, O.O., Bengtsson, A. and Erlinge, D. 2008. Gene expression profiling for the identification of G protein-coupled receptors in human platelets. *Thromb. Res.* 122: 47-57.

CHROMOSOMAL LOCATION

Genetic locus: GPR101 (human) mapping to Xq26.3.

PRODUCT

GPR101 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see GPR101 shRNA Plasmid (h): sc-75166-SH and GPR101 shRNA (h) Lentiviral Particles: sc-75166-V as alternate gene silencing products.

For independent verification of GPR101 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-75166A, sc-75166B and sc-75166C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

GPR101 siRNA (h) is recommended for the inhibition of GPR101 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GPR101 gene expression knockdown using RT-PCR Primer: GPR101 (h)-PR: sc-75166-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.