

GPR19 siRNA (m): sc-75174

BACKGROUND

G protein-coupled receptors (GPRs or GPCRs), also known as seven transmembrane receptors, heptahelical receptors, or 7TM receptors, are members of the largest protein family and play a role in many different stimulus-response pathways. G protein-coupled receptors mediate extracellular signals into intracellular signals (G-protein activation). They respond to a great variety of signaling molecules, including hormones, neurotransmitters and other proteins and peptides. GPR proteins are integral seven-pass membrane proteins with some conserved amino acid regions. GPR19, an orphan receptor, shows elevated expression during embryonic development of the nervous system as well as in specific regions of adult mouse brain, including the olfactory bulb, the hippocampus, hypothalamic nuclei and the cerebellum. The GPR19 gene maps to a location on chromosome 12, which is a frequent target for rearrangement in cancer cells and involved in childhood acute lymphoblastic leukemia (ALL).

REFERENCES

1. O'Dowd, B.F., Nguyen, T., Lynch, K.R., Kolakowski, L.F., Thompson, M., Cheng, R., Marchese, A., Ng, G., Heng, H.H. and George, S.R. 1996. A novel gene codes for a putative G protein-coupled receptor with an abundant expression in brain. *FEBS Lett.* 394: 325-329.
2. Montpetit, A. and Sinnett, D. 1999. Physical mapping of the G protein-coupled receptor 19 (GPR19) in the chromosome 12p12.3 region frequently rearranged in cancer cells. *Hum. Genet.* 105: 162-164.
3. Bresnick, J.N., Skynner, H.A., Chapman, K.L., Jack, A.D., Zamiara, E., Negulescu, P., Beaumont, K., Patel, S. and McAllister, G. 2003. Identification of signal transduction pathways used by orphan G protein-coupled receptors. *Assay Drug Dev. Technol.* 1: 239-249.
4. Hoffmeister-Ullrich, S.A., Süsens, U. and Schaller, H.C. 2004. The orphan G protein-coupled receptor GPR19 is expressed predominantly in neuronal cells during mouse embryogenesis. *Cell Tissue Res.* 318: 459-463.
5. Rossi, P., Dolci, S., Sette, C., Capolunghi, F., Pellegrini, M., Loiarro, M., Di Agostino, S., Paronetto, M.P., Grimaldi, P., Merico, D., Martegani, E. and Geremia, R. 2004. Analysis of the gene expression profile of mouse male meiotic germ cells. *Gene Expr. Patterns* 4: 267-281.
6. Assou, S., Le Carrou, T., Tondeur, S., Ström, S., Gabelle, A., Marty, S., Nadal, L., Pantesco, V., Reme, T., Hugnot, J.P., Gasca, S., Hovatta, O., Hamamah, S., Klein, B. and De Vos, J. 2007. A meta-analysis of human embryonic stem cells transcriptome integrated into a web-based expression atlas. *Stem Cells* 25: 961-973.
7. Riker, A.I., Enkemann, S.A., Fodstad, O., Liu, S., Ren, S., Morris, C., Xi, Y., Howell, P., Metge, B., Samant, R.S., Shevde, L.A., Li, W., Eschrich, S., Daud, A., Ju, J. and Matta, J. 2008. The gene expression profiles of primary and metastatic melanoma yields a transition point of tumor progression and metastasis. *BMC Med. Genomics* 1: 13.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: Gpr19 (mouse) mapping to 6 G1.

PRODUCT

GPR19 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see GPR19 shRNA Plasmid (m): sc-75174-SH and GPR19 shRNA (m) Lentiviral Particles: sc-75174-V as alternate gene silencing products.

For independent verification of GPR19 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-75174A, sc-75174B and sc-75174C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

GPR19 siRNA (m) is recommended for the inhibition of GPR19 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GPR19 gene expression knockdown using RT-PCR Primer: GPR19 (m)-PR: sc-75174-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.