

GRTP1 siRNA (h): sc-75204

BACKGROUND

Pituitary growth hormone (GH), also designated somatotropin, plays a crucial role in stimulating and controlling the growth, metabolism and differentiation of many mammalian cell types by modulating the synthesis of multiple mRNA species. These effects are mediated by the binding of GH to its membrane-bound receptor, GHR, and involve a phosphorylation cascade that results in the modulation of numerous signaling pathways. GRTP1 (growth hormone-regulated TBC protein 1), also designated TBC1 domain family member six, is a 336 amino acid protein with expression modulated by growth hormone. The TBC1 domain of GRTP1 is a region that is notably conserved among GTPase-activating proteins, which characteristically play a role in the cell cycle and mitosis coordination. Interestingly, expression of GRTP1 is significantly upregulated in post-puberty testis and then maintained in the adult, suggesting that the germ cell may be the major site of GRTP1 function. There are two isoforms of GRTP1, which are a result of alternative splicing events.

REFERENCES

1. Lobie, P.E., Breipohl, W., Aragón, J.G. and Waters, M.J. 1990. Cellular localization of the growth hormone receptor/binding protein in the male and female reproductive systems. *Endocrinology* 126: 2214-2221.
2. Boguski, M.S. and McCormick, F. 1993. Proteins regulating Ras and its relatives. *Nature* 366: 643-654.
3. Gruaz, N.M., Arsenijevic, Y., Wehrenberg, W.B., Sizonenko, P.C. and Aubert, M.L. 1994. Growth hormone (GH) deprivation induced by passive immunization against rat GH-releasing factor does not disturb the course of sexual maturation and fertility in the female rat. *Endocrinology* 135: 509-519.
4. Breier, B.H., Vickers, M.H., Gravance, C.G. and Casey, P.J. 1996. Growth hormone (GH) therapy markedly increases the motility of spermatozoa and the concentration of Insulin-like growth factor-I in seminal vesicle fluid in the male GH-deficient dwarf rat. *Endocrinology* 137: 4061-4064.
5. Lu, C., Kasik, J., Stephan, D.A., Yang, S., Sperling, M.A. and Menon, R.K. 2001. GRTP1, a novel gene regulated by growth hormone. *Endocrinology* 142: 4568-4571.
6. Pan, X., Eathiraj, S., Munson, M. and Lambright, D.G. 2006. TBC-domain GAPs for Rab GTPases accelerate GTP hydrolysis by a dual-finger mechanism. *Nature* 442: 303-306.

CHROMOSOMAL LOCATION

Genetic locus: GRTP1 (human) mapping to 13q34.

PRODUCT

GRTP1 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see GRTP1 shRNA Plasmid (h): sc-75204-SH and GRTP1 shRNA (h) Lentiviral Particles: sc-75204-V as alternate gene silencing products.

For independent verification of GRTP1 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-75204A, sc-75204B and sc-75204C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

GRTP1 siRNA (h) is recommended for the inhibition of GRTP1 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

GRTP1 (G-6): sc-398974 is recommended as a control antibody for monitoring of GRTP1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GRTP1 gene expression knockdown using RT-PCR Primer: GRTP1 (h)-PR: sc-75204-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.