



# HELIC1 siRNA (h): sc-75241

## BACKGROUND

HELIC1 (helicase, ATP binding 1), also known as ASCC3 (activating signal co-integrator 1 complex subunit 3) or RNAH, is a 2,202 amino acid protein that localizes to the cytoplasm and contains two helicase ATP-binding domains, two helicase C-terminal domains and three SEC63 domains. Expressed ubiquitously, HELIC1 exists as a component of the multi-protein TRIP4 complex and is thought to enhance the transactivation of NFκB, SRF and c-Jun. The gene encoding HELIC1 maps to human chromosome 6, which contains 170 million base pairs and comprises nearly 6% of the human genome. Deletion of a portion of the q arm of chromosome 6 is associated with early onset intestinal cancer, suggesting the presence of a cancer susceptibility locus. Additionally, Porphyria cutanea tarda, Parkinson's disease, Stickler syndrome and a susceptibility to bipolar disorder are all associated with genes that map to chromosome 6.

## REFERENCES

1. Coulie, P.G., et al. 1999. Antitumor immunity at work in a melanoma patient. *Adv. Cancer Res.* 76: 213-242.
2. Makino, N., et al. 2001. Isolation and characterization of the human gene homologous to the *Drosophila* headcase (hdc) gene in chromosome bands 6q23-q24, a region of common deletion in human pancreatic cancer. *DNA Seq.* 11: 547-553.
3. Jung, D.J., et al. 2002. Novel transcription coactivator complex containing activating signal co-integrator 1. *Mol. Cell. Biol.* 22: 5203-5211.
4. McQueen, M.B., et al. 2005. Combined analysis from eleven linkage studies of bipolar disorder provides strong evidence of susceptibility loci on chromosomes 6q and 8q. *Am. J. Hum. Genet.* 77: 582-595.
5. Chien, C.C., et al. 2006. A homologue of the *Drosophila* headcase protein is a novel tumor marker for early-stage colorectal cancer. *Oncol. Rep.* 15: 919-926.
6. Di Meo, G.P., et al. 2007. An advanced sheep (*Ovis aries*, 2n=54) cytogenetic map and assignment of 88 new autosomal loci by fluorescence *in situ* hybridization and R-banding. *Anim. Genet.* 38: 233-240.

## CHROMOSOMAL LOCATION

Genetic locus: ASCC3 (human) mapping to 6q16.3.

## PRODUCT

HELIC1 siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see HELIC1 shRNA Plasmid (h): sc-75241-SH and HELIC1 shRNA (h) Lentiviral Particles: sc-75241-V as alternate gene silencing products.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

HELIC1 siRNA (h) is recommended for the inhibition of HELIC1 expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor HELIC1 gene expression knockdown using RT-PCR Primer: HELIC1 (h)-PR: sc-75241-PR (20 μl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.