## SANTA CRUZ BIOTECHNOLOGY, INC.

# Josephin-1 siRNA (h): sc-75361



## BACKGROUND

Josephin-1, also known as JOSD1 (josephin domain containing 1), is a 202 amino acid protein that contains one josephin domain and is encoded by a gene that maps to human chromosome 22. Chromosome 22 houses over 500 genes and is the second smallest human chromosome. Mutations in several of the genes that map to chromosome 22 are involved in the development of Phelan-McDermid syndrome, neurofibromatosis type 2, autism and schizophrenia. Additionally, translocations between chromosomes 9 and 22 may lead to the formation of the Philadelphia chromosome and the subsequent production of the novel fusion protein Bcr-Abl, a potent cell proliferation activator found in several types of leukemias.

## REFERENCES

- Gilbert, F. 1998. Disease genes and chromosomes: disease maps of the human genome. Chromosome 22. Genet. Test. 2: 89-97.
- 2. Schwab, S.G. and Wildenauer, D.B. 1999. Chromosome 22 workshop report. Am. J. Med. Genet. 88: 276-278.
- 3. Matsuda, A., et al. 2003. Large-scale identification and characterization of human genes that activate NF $\kappa$ B and MAPK signaling pathways. Oncogene 22: 3307-3318.
- Arinami, T. 2006. Analyses of the associations between the genes of 22q11 deletion syndrome and schizophrenia. J. Hum. Genet. 51: 1037-1045.
- Paylor, R., et al. 2006. Tbx1 haploinsufficiency is linked to behavioral disorders in mice and humans: implications for 22q11 deletion syndrome. Proc. Natl. Acad. Sci. USA 103: 7729-7734.
- Zheng, X., et al. 2006. Bcr and its mutants, the reciprocal t(9;22)-associated Abl/Bcr fusion proteins, differentially regulate the cytoskeleton and cell motility. BMC Cancer 6: 262.

## CHROMOSOMAL LOCATION

Genetic locus: JOSD1 (human) mapping to 22q13.1.

## PRODUCT

Josephin-1 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Josephin-1 shRNA Plasmid (h): sc-75361-SH and Josephin-1 shRNA (h) Lentiviral Particles: sc-75361-V as alternate gene silencing products.

For independent verification of Josephin-1 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-75361A, sc-75361B and sc-75361C.

#### **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

#### PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

#### STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNAse-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNAse-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## **APPLICATIONS**

Josephin-1 siRNA (h) is recommended for the inhibition of Josephin-1 expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

### **RT-PCR REAGENTS**

Semi-quantitative RT-PCR may be performed to monitor Josephin-1 gene expression knockdown using RT-PCR Primer: Josephin-1 (h)-PR: sc-75361-PR (20  $\mu$ I). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.