

LPLUNC4 siRNA (h): sc-75696

BACKGROUND

LPLUNC4 (long palate, lung and nasal epithelium carcinoma-associated protein 4), also known as C20orf186 or RY2G5, is a 614 amino acid secreted protein that belongs to the BPI/LBP/PLUNC superfamily. Expressed in nasal tissue, LPLUNC4 has the capacity to recognize and bind specific classes of odorants serving as a primary defense mechanism in the mucosa. LPLUNC4 is encoded by a gene which maps to a BPI/LBP/PLUNC gene cluster on human chromosome 20q11.21. Comprising approximately 2% of the human genome, chromosome 20 contains nearly 63 million bases that encode over 600 genes, some of which are associated with Creutzfeldt-Jakob disease, amyotrophic lateral sclerosis, spinal muscular atrophy, ring chromosome 20 epilepsy syndrome and Alagille syndrome.

REFERENCES

1. Bingle, C.D. and Craven, C.J. 2002. PLUNC: a novel family of candidate host defence proteins expressed in the upper airways and nasopharynx. *Hum. Mol. Genet.* 11: 937-943.
2. Zhang, B., Nie, X., Xiao, B., Xiang, J., Shen, S., Gong, J., Zhou, M., Zhu, S., Zhou, J., Qian, J., Lu, H., He, X., Li, X., Hu, G. and Li, G. 2003. Identification of tissue-specific genes in nasopharyngeal epithelial tissue and differentially expressed genes in nasopharyngeal carcinoma by suppression subtractive hybridization and cDNA microarray. *Genes Chromosomes Cancer* 38: 80-90.
3. Hou, J., Yashiro, K., Okazaki, Y., Saijoh, Y., Hayashizaki, Y. and Hamada, H. 2004. Identification of a novel left-right asymmetrically expressed gene in the mouse belonging to the BPI/PLUNC superfamily. *Dev. Dyn.* 229: 373-379.
4. LeClair, E.E., Nomellini, V., Bahena, M., Singleton, V., Bingle, L., Craven, C.J. and Bingle, C.D. 2004. Cloning and expression of a mouse member of the PLUNC protein family exclusively expressed in tongue epithelium. *Genomics* 83: 658-666.
5. Casado, B., Pannell, L.K., Iadarola, P. and Baraniuk, J.N. 2005. Identification of human nasal mucous proteins using proteomics. *Proteomics* 5: 2949-2959.
6. Flach, C.F., Qadri, F., Bhuiyan, T.R., Alam, N.H., Jennische, E., Lönnroth, I. and Holmgren, J. 2007. Broad upregulation of innate defense factors during acute cholera. *Infect. Immun.* 75: 2343-2350.
7. Vargas, P.A., Speight, P.M., Bingle, C.D., Barrett, A.W. and Bingle, L. 2008. Expression of PLUNC family members in benign and malignant salivary gland tumours. *Oral Dis.* 14: 613-619.

CHROMOSOMAL LOCATION

Genetic locus: BPIFB4 (human) mapping to 20q11.21.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

LPLUNC4 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see LPLUNC4 shRNA Plasmid (h): sc-75696-SH and LPLUNC4 shRNA (h) Lentiviral Particles: sc-75696-V as alternate gene silencing products.

For independent verification of LPLUNC4 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-75696A, sc-75696B and sc-75696C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

LPLUNC4 siRNA (h) is recommended for the inhibition of LPLUNC4 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor LPLUNC4 gene expression knockdown using RT-PCR Primer: LPLUNC4 (h)-PR: sc-75696-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.