

Myozenin 3 siRNA (m): sc-75855

BACKGROUND

Myozenins, which includes Myozenin 1, Myozenin 2 and Myozenin 3, are a family of intracellular binding proteins that are involved in linking Z-disk proteins to the sarcomere. Myozenin 3, also known as calsarcin-3 or FATZ-related protein 3 (FRP3), is a 251 amino acid protein expressed specifically in skeletal muscle and enriched in fast-twitch muscle fibers. Myozenin 3, like Myozenin 1 and 2, interacts with calnecurin and the Z-disc proteins α -actinin, Filamin 2, and Telethonin. Myozenin 3 plays an important role in the modulation of calnecurin signaling, and it is thought to play a role in myofibrillogenesis. Due to the close involvement of Myozenin 3 with muscle formation, mutations in the gene encoding Myozenin 3 may be associated with muscular dystrophies and neuromuscular myopathies. Three isoforms of Myozenin 3 exist as a result of alternative splicing events.

REFERENCES

1. Frey, N. and Olson, E.N. 2002. Calsarcin-3, a novel skeletal muscle-specific member of the calsarcin family, interacts with multiple Z-disc proteins. *J. Biol. Chem.* 277: 13998-14004.
2. Martin, L.J., Comuzzie, A.G., Sonnenberg, G.E., Myklebust, J., James, R., Marks, J., Blangero, J. and Kissebah, A.H. 2004. Major quantitative trait locus for resting heart rate maps to a region on chromosome 4. *Hypertension* 43: 1146-1151.
3. Gontier, Y., Taivainen, A., Fontao, L., Sonnenberg, A., van der Flier, A., Carpen, O., Faulkner, G. and Borradori, L. 2005. The Z-disc proteins myotilin and FATZ-1 interact with each other and are connected to the sarcolemma via muscle-specific filamins. *J. Cell Sci.* 118: 3739-3749.
4. Wang, H., Yang, S., Yang, E., Zhu, Z., Mu, Y., Feng, S. and Li, K. 2007. NF κ B mediates the transcription of mouse calsarcin-1 gene, but not calsarcin-2, in C2C12 cells. *BMC Mol. Biol.* 8: 19.
5. Schoensiegel, F., Bekeredian, R., Schrewe, A., Weichenhan, D., Frey, N., Katus, H.A. and Ivandic, B.T. 2007. Atrial natriuretic peptide and osteopontin are useful markers of cardiac disorders in mice. *Comp. Med.* 57: 546-553.
6. Arola, A.M., Sanchez, X., Murphy, R.T., Hasle, E., Li, H., Elliott, P.M., McKenna, W.J., Towbin, J.A. and Bowles, N.E. 2007. Mutations in PDLIM3 and MYOZ1 encoding myocyte Z line proteins are infrequently found in idiopathic dilated cardiomyopathy. *Mol. Genet. Metab.* 90: 435-440.
7. Online Mendelian Inheritance in Man, OMIM[™]. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 610735. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
8. von Nandelstadh, P., Ismail, M., Gardin, C., Suila, H., Zara, I., Belgrano, A., Valle, G., Carpen, O. and Faulkner, G. 2009. A class III PDZ binding motif in the myotilin and FATZ families binds enigma family proteins: a common link for Z-disc myopathies. *Mol. Cell. Biol.* 29: 822-834.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: Myoz3 (mouse) mapping to 18 D3.

PRODUCT

Myozenin 3 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Myozenin 3 shRNA Plasmid (m): sc-75855-SH and Myozenin 3 shRNA (m) Lentiviral Particles: sc-75855-V as alternate gene silencing products.

For independent verification of Myozenin 3 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-75855A, sc-75855B and sc-75855C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Myozenin 3 siRNA (m) is recommended for the inhibition of Myozenin 3 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Myozenin 3 gene expression knockdown using RT-PCR Primer: Myozenin 3 (m)-PR: sc-75855-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.