

NOL4 siRNA (h): sc-75943

BACKGROUND

The nucleolus consists of a number of specific proteins that play a critical role in the assembly of ribosomes, as well as in the maintenance and structural integrity of the nucleolus. NOL4 (nucleolar protein 4), also known as NOLP (nucleolar-localized protein), is a 524 amino acid protein that is predominantly expressed in testis and brain. NOL4 contains at least two domains that direct it to subnuclear locations. The gene encoding NOL4 is located on human chromosome 18, which houses over 300 protein-coding genes and contains nearly 76 million bases. There are a variety of diseases associated with defects in chromosome 18-localized genes, some of which include Trisomy 18 (also known as Edwards syndrome), Niemann-Pick disease, hereditary hemorrhagic telangiectasia, erythropoietic protoporphyria and follicular lymphomas. There are two isoforms of NOL4 that are produced as a result of alternative splicing events.

REFERENCES

1. Carstea, E.D., Polymeropoulos, M.H., Parker, C.C., Detera-Wadleigh, S.D., O'Neill, R.R., Patterson, M.C., Goldin, E., Xiao, H., Straub, R.E. and Vanier, M.T. 1993. Linkage of Niemann-Pick disease type C to human chromosome 18. *Proc. Natl. Acad. Sci. USA* 90: 2002-2004.
2. Ueki, N., Kondo, M., Seki, N., Yano, K., Oda, T., Masuho, Y. and Muramatsu, M. 1998. NOLP: identification of a novel human nucleolar protein and determination of sequence requirements for its nucleolar localization. *Biochem. Biophys. Res. Commun.* 252: 97-102.
3. Ueki, N., Oda, T., Kondo, M., Yano, K., Noguchi, T. and Muramatsu, M. 1998. Selection system for genes encoding nuclear-targeted proteins. *Nat. Biotechnol.* 16: 1338-1342.
4. Zhu, L., Swergold, G.D. and Seldin, M.F. 2003. Examination of sequence homology between human chromosome 20 and the mouse genome: intense conservation of many genomic elements. *Hum. Genet.* 113: 60-70.
5. Online Mendelian Inheritance in Man, OMIM[™]. 2003. Johns Hopkins University, Baltimore, MD. MIM Number: 603577. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Grosso, S., Pucci, L., Di Bartolo, R.M., Gobbi, G., Bartalini, G., Anichini, C., Scarinci, R., Balestri, M., Farnetani, M.A., Cioni, M., Morgese, G. and Balestri, P. 2005. Chromosome 18 aberrations and epilepsy: a review. *Am. J. Med. Genet. A* 134A: 88-94.
7. Wang, S.S., Smiraglia, D.J., Wu, Y.Z., Ghosh, S., Rader, J.S., Cho, K.R., Bonfiglio, T.A., Nayar, R., Plass, C. and Sherman, M.E. 2008. Identification of novel methylation markers in cervical cancer using restriction landmark genomic scanning. *Cancer Res.* 68: 2489-2497.
8. Cerminara, C., Lo Castro, A., D'Argenzio, L., Galasso, C. and Curatolo, P. 2008. Epilepsy and deletion syndromes of chromosome 18: do not forget the short arm! *Epilepsia* 49: 1813-1814.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: NOL4 (human) mapping to 18q12.1.

PRODUCT

NOL4 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see NOL4 shRNA Plasmid (h): sc-75943-SH and NOL4 shRNA (h) Lentiviral Particles: sc-75943-V as alternate gene silencing products.

For independent verification of NOL4 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-75943A, sc-75943B and sc-75943C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

NOL4 siRNA (h) is recommended for the inhibition of NOL4 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor NOL4 gene expression knockdown using RT-PCR Primer: NOL4 (h)-PR: sc-75943-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.