

GluR-3 (C-20): sc-7612

BACKGROUND

Glutamate receptors mediate most excitatory neurotransmission in the brain and play an important role in neural plasticity, neural development and neurodegeneration. Ionotropic glutamate receptors are categorized into NMDA receptors and kainate/AMPA receptors, both of which contain glutamate-gated, cation-specific ion channels. Kainate/AMPA receptors are co-localized with NMDA receptors in many synapses and consist of seven structurally related subunits designated GluR-1 to -7. The kainate/AMPA receptors are primarily responsible for the fast excitatory neurotransmission by glutamate, whereas the NMDA receptors are functionally characterized by a slow kinetic and a high permeability for Ca^{2+} ions. The NMDA receptors consist of five subunits: ϵ 1, 2, 3, 4 and one ζ subunit. The ζ subunit is expressed throughout the brainstem, whereas the four ϵ subunits display limited distribution.

REFERENCES

1. Choi, D.W., et al. 1990. The role of glutamate neurotoxicity in hypoxic-ischemic neuronal death. *Annu. Rev. Neurosci.* 13: 171-182.
2. Stern, P., et al. 1992. Fast and slow components of unitary EPSCs on stellate cells elicited by focal stimulation in slices of rat visual cortex. *J. Physiol.* 449: 247-278.

CHROMOSOMAL LOCATION

Genetic locus: GRIA3 (human) mapping to Xq25, GRIA2 (human) mapping to 4q32.1; Gria3 (mouse) mapping to X A3.3, Gria2 (mouse) mapping to 3 E3.

SOURCE

GluR-3 (C-20) is an affinity purified goat polyclonal antibody raised against a peptide mapping at the C-terminus of GluR-3 of human origin.

PRODUCT

Each vial contains 200 μg IgG in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

Blocking peptide available for competition studies, sc-7612 P, (100 μg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% BSA).

APPLICATIONS

GluR-3 (C-20) is recommended for detection of GluR-3 and, to a lesser extent, GluR-2 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 μg per 100-500 μg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).

GluR-3 (C-20) is also recommended for detection of GluR-3 and, to a lesser extent, GluR-2 in additional species, including equine, canine, bovine, porcine and avian.


Molecular Weight of GluR-3: 103 kDa.

Positive Controls: mouse brain extract: sc-2253, rat brain extract: sc-2392 or mouse cerebellum extract: sc-2403.


RECOMMENDED SECONDARY REAGENTS

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker[™] Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunoprecipitation: use Protein A/G PLUS-Agarose: sc-2003 (0.5 ml agarose/2.0 ml). 3) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

DATA



GluR-3 (C-20): sc-7612. Western blot analysis of GluR-3 expression in rat brain (A) and mouse brain (B) tissue extracts.



GluR-3 (C-20): sc-7612. Western blot analysis of GluR-3 expression in mouse cerebellum tissue extract.

SELECT PRODUCT CITATIONS

1. Puller, C., et al. 2011. Cell-type-specific localization of protocadherin $\beta 16$ at AMPA and AMPA/Kainate receptor-containing synapses in the primate retina. *J. Comp. Neurol.* 519: 467-479.

STORAGE

Store at 4° C, **DO NOT FREEZE**. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

MONOS
Satisfaction
Guaranteed

Try **GluR-3 (1D2E2): sc-517202**, our highly recommended monoclonal alternative to GluR-3 (C-20).