

POLDIP2 siRNA (h): sc-76190

BACKGROUND

POLDIP2 (polymerase (DNA-directed), δ interacting protein 2), also known as POLD4 or PDIP38, is a 368 amino acid protein that localizes to the nucleus and contains one apaG domain. Interacting with PCNA and DNA pol δ 2, POLDIP2 is thought to influence DNA replication and cellular proliferation events, specifically by inhibiting the activity of DNA pol subunits. Human POLDIP2 shares 95% sequence identity with its mouse counterpart, suggesting a conserved role between species. The gene encoding POLDIP2 maps to human chromosome 17, which comprises over 2.5% of the human genome and encodes over 1,200 genes. Two key tumor suppressor genes are associated with chromosome 17, namely, p53 and BRCA1. Tumor suppressor p53 is necessary for maintenance of cellular genetic integrity by moderating cell fate through DNA repair versus cell death. Malfunction or loss of p53 expression is associated with malignant cell growth and Li-Fraumeni syndrome. Like p53, BRCA1 is directly involved in DNA repair, though specifically it is recognized as a genetic determinant of early onset breast cancer and predisposition to cancers of the ovary, colon, prostate gland and fallopian tubes.

REFERENCES

- Hartley, J.L., Temple, G.F. and Brasch, M.A. 2000. DNA cloning using *in vitro* site-specific recombination. *Genome Res.* 10: 1788-1795.
- Liu, L., Rodriguez-Belmonte, E.M., Mazloum, N., Xie, B. and Lee, M.Y. 2003. Identification of a novel protein, PDIP38, that interacts with the p50 subunit of DNA polymerase δ and proliferating cell nuclear antigen. *J. Biol. Chem.* 278: 10041-10047.
- Xie, B., Li, H., Wang, Q., Xie, S., Rahmeh, A., Dai, W. and Lee, M.Y. 2005. Further characterization of human DNA polymerase δ interacting protein 38. *J. Biol. Chem.* 280: 22375-22384.
- Klaile, E., Müller, M.M., Kannicht, C., Otto, W., Singer, B.B., Reutter, W., Obrink, B. and Lucka, L. 2007. The cell adhesion receptor carcinoembryonic antigen-related cell adhesion molecule 1 regulates nucleocytoplasmic trafficking of DNA polymerase δ -interacting protein 38. *J. Biol. Chem.* 282: 26629-26640.

CHROMOSOMAL LOCATION

Genetic locus: POLDIP2 (human) mapping to 17q11.2.

PRODUCT

POLDIP2 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see POLDIP2 shRNA Plasmid (h): sc-76190-SH and POLDIP2 shRNA (h) Lentiviral Particles: sc-76190-V as alternate gene silencing products.

For independent verification of POLDIP2 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-76190A, sc-76190B and sc-76190C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

POLDIP2 siRNA (h) is recommended for the inhibition of POLDIP2 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor POLDIP2 gene expression knockdown using RT-PCR Primer: POLDIP2 (h)-PR: sc-76190-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.