



Rab 19 siRNA (h): sc-76318

BACKGROUND

The Ras-related superfamily of guanine nucleotide binding proteins includes the R-Ras, Rap, Ral/Rec and Rho/Rab subfamilies, all of which are thought to play an important role in either endocytosis or in biosynthetic protein transport. The process of transporting newly synthesized proteins from the endoplasmic reticulum (ER) to various stacks of the Golgi complex and to secretory vesicles involves the movement of carrier vesicles and requires Rab protein function. Rab proteins are also an integral part of endocytic pathways. Rab 19 is a 217 amino acid protein that is lipid anchored to the cytoplasmic side of the cell membrane and, as a member of the small GTPase superfamily, may play a role in endocytotic pathways. Multiple isoforms of Rab 19 exist due to alternative splicing events.

REFERENCES

1. Olkkonen, V.M., Dupree, P., Killisch, I., Lütcke, A., Zerial, M. and Simons, K. 1993. Molecular cloning and subcellular localization of three GTP-binding proteins of the rab subfamily. *J. Cell Sci.* 106: 1249-1261.
2. Chen, D., Guo, J. and Gahl, W.A. 1997. RAB GTPases expressed in human melanoma cells. *Biochim. Biophys. Acta* 1355: 1-6.
3. Zhao, H., Ettala, O. and Väänänen, H.K. 2002. Intracellular membrane trafficking pathways in bone-resorbing osteoclasts revealed by cloning and subcellular localization studies of small GTP-binding rab proteins. *Biochem. Biophys. Res. Commun.* 293: 1060-1065.
4. Ali, B.R., Wasmeier, C., Lamoreux, L., Strom, M. and Seabra, M.C. 2004. Multiple regions contribute to membrane targeting of Rab GTPases. *J. Cell Sci.* 117: 6401-6412.
5. Chakrabarty, K. and Heumann, R. 2008. Prospective of Ras signaling in stem cells. *Biol. Chem.* 389: 791-798.
6. Fukuda, M., Kanno, E., Ishibashi, K. and Itoh, T. 2008. Large scale screening for novel rab effectors reveals unexpected broad Rab binding specificity. *Mol. Cell. Proteomics* 7: 1031-1042.

CHROMOSOMAL LOCATION

Genetic locus: RAB19 (human) mapping to 7q34.

PRODUCT

Rab 19 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Rab 19 shRNA Plasmid (h): sc-76318-SH and Rab 19 shRNA (h) Lentiviral Particles: sc-76318-V as alternate gene silencing products.

For independent verification of Rab 19 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-76318A, sc-76318B and sc-76318C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Rab 19 siRNA (h) is recommended for the inhibition of Rab 19 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Rab 19 gene expression knockdown using RT-PCR Primer: Rab 19 (h)-PR: sc-76318-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.