

RASL10B siRNA (h): sc-76353

BACKGROUND

Ras-like protein family member 10B (RASL10B), also known as Ras-like protein VTS58635, is a 203 amino acid protein that belongs to the Ras family. Localized to the cytoplasmic side of the cell membrane, RASL10B is widely expressed in human tissues and is a candidate tumor suppressor protein. The gene encoding RASL10B maps to chromosome 17q12. Chromosome 17 makes up over 2.5% of the human genome with about 81 million bases encoding over 1,200 genes. Two key tumor suppressor genes are associated with chromosome 17, namely, p53 and BRCA1. Chromosome 17 is also linked to neurofibromatosis, a condition characterized by neural and epidermal lesions, and dysregulated Schwann cell growth. Alexander disease, Birt-Hogg-Dube syndrome and Canavan disease are also associated with chromosome 17.

REFERENCES

1. Bos, J.L. 1997. Ras-like GTPases. *Biochim. Biophys. Acta* 1333: M19-M31.
2. Ellis, C.A., Vos, M.D., Howell, H., Vallecorsa, T., Fuets, D.W. and Clark, G.J. 2002. Rig is a novel Ras-related protein and potential neural tumor suppressor. *Proc. Natl. Acad. Sci. USA* 99: 9876-9881.
3. Zou, H., Hu, L., Li, J., Zhan, S. and Cao, K. 2006. Cloning and characterization of a novel small monomeric GTPase, RASL10B, with tumor suppressor potential. *Biotechnol. Lett.* 28: 1901-1908.
4. Tuskan, R.G., Tsang, S., Sun, Z., Baer, J., Rozenblum, E., Wu, X., Munroe, D.J. and Reilly, K.M. 2008. Real-time PCR analysis of candidate imprinted genes on mouse chromosome 11 shows balanced expression from the maternal and paternal chromosomes and strain-specific variation in expression levels. *Epigenetics* 3: 43-50.
5. Neuwald, A.F. 2009. The glycine brace: a component of Rab, Rho, and Ran GTPases associated with hinge regions of guanine- and phosphate-binding loops. *BMC Struct. Biol.* 9: 11.
6. Pannekoek, W.J., Kooistra, M.R., Zwartkruis, F.J. and Bos, J.L. 2009. Cell-cell junction formation: the role of Rap1 and Rap1 guanine nucleotide exchange factors. *Biochim. Biophys. Acta* 1788: 790-796.

CHROMOSOMAL LOCATION

Genetic locus: RASL10B (human) mapping to 17q12.

PRODUCT

RASL10B siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see RASL10B shRNA Plasmid (h): sc-76353-SH and RASL10B shRNA (h) Lentiviral Particles: sc-76353-V as alternate gene silencing products.

For independent verification of RASL10B (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-76353A, sc-76353B and sc-76353C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

RASL10B siRNA (h) is recommended for the inhibition of RASL10B expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor RASL10B gene expression knockdown using RT-PCR Primer: RASL10B (h)-PR: sc-76353-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.