

# SELB siRNA (h): sc-76468

## BACKGROUND

SELB (selenocysteine-specific elongation factor), also known as EEFSEC (eukaryotic elongation factor, selenocysteine-tRNA-specific) or EFSEC, is a 596 amino acid protein that localizes to both the nucleus and the cytoplasm and belongs to the GTP-binding elongation factor family. Functioning as a translation factor, SELB binds GTP and GDP and is necessary for the incorporation of selenocysteine into target proteins. The gene encoding SELB maps to human chromosome 3, which houses over 1,100 genes, including a chemokine receptor (CKR) gene cluster and a variety of human cancer-related gene loci. Key tumor suppressing genes on chromosome 3 include those that encode the apoptosis mediator RASSF1, the cell migration regulator HYAL1 and the angiogenesis suppressor SEMA3B. Marfan syndrome, porphyria, von Hippel-Lindau syndrome, osteogenesis imperfecta and Charcot-Marie-Tooth disease are a few of the numerous genetic diseases associated with chromosome 3.

## REFERENCES

1. Fagegaltier, D., et al. 2000. Characterization of mSelB, a novel mammalian elongation factor for selenoprotein translation. *EMBO J.* 19: 4796-4805.
2. Zavacki, A.M., et al. 2003. Coupled tRNA(Sec)-dependent assembly of the selenocysteine decoding apparatus. *Mol. Cell* 11: 773-781.
3. Online Mendelian Inheritance in Man, OMIM™. 2003. Johns Hopkins University, Baltimore, MD. MIM Number: 607695. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Mehta, A., et al. 2004. Efficiency of mammalian selenocysteine incorporation. *J. Biol. Chem.* 279: 37852-37859.
5. Caban, K. and Copeland, P.R. 2006. Size matters: a view of selenocysteine incorporation from the ribosome. *Cell. Mol. Life Sci.* 63: 73-81.
6. Gupta, M. and Copeland, P.R. 2007. Functional analysis of the interplay between translation termination, selenocysteine codon context, and selenocysteine insertion sequence-binding protein 2. *J. Biol. Chem.* 282: 36797-36807.
7. Caban, K., et al. 2007. The L7Ae RNA binding motif is a multifunctional domain required for the ribosome-dependent Sec incorporation activity of Sec insertion sequence binding protein 2. *Mol. Cell. Biol.* 27: 6350-6360.

## CHROMOSOMAL LOCATION

Genetic locus: EEFSEC (human) mapping to 3q21.3.

## PRODUCT

SELB siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SELB shRNA Plasmid (h): sc-76468-SH and SELB shRNA (h) Lentiviral Particles: sc-76468-V as alternate gene silencing products.

For independent verification of SELB (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-76468A, sc-76468B and sc-76468C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

SELB siRNA (h) is recommended for the inhibition of SELB expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

SELB (G-9): sc-166521 is recommended as a control antibody for monitoring of SELB gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SELB gene expression knockdown using RT-PCR Primer: SELB (h)-PR: sc-76468-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.