



SOHLH1 siRNA (h): sc-76537

BACKGROUND

SOHLH1 (spermatogenesis and oogenesis specific basic helix-loop-helix 1), also known as NOHLH or TEB2, is a 328 amino acid protein that localizes to both the nucleus and the cytoplasm and contains one bHLH domain through which it may function as a transcription factor during oogenesis and spermatogenesis. The gene encoding SOHLH1 maps to human chromosome 9, which houses over 900 genes and comprises nearly 4% of the human genome. Hereditary hemorrhagic telangiectasia, which is characterized by harmful vascular defects, and Familial dysautonomia, are both associated with chromosome 9. Notably, chromosome 9 encompasses the largest interferon family gene cluster.

REFERENCES

1. Ballow, D., Meistrich, M.L., Matzuk, M. and Rajkovic, A. 2006. SOHLH1 is essential for spermatogonial differentiation. *Dev. Biol.* 294: 161-167.
2. Pangas, S.A., Choi, Y., Ballow, D.J., Zhao, Y., Westphal, H., Matzuk, M.M. and Rajkovic, A. 2006. Oogenesis requires germ cell-specific transcriptional regulators SOHLH1 and Lhx8. *Proc. Natl. Acad. Sci. USA* 103: 8090-8095.
3. Suzumori, N., Pangas, S.A. and Rajkovic, A. 2007. Candidate genes for premature ovarian failure. *Curr. Med. Chem.* 14: 353-357.
4. Dadoune, J.P. 2007. New insights into male gametogenesis: what about the spermatogonial stem cell niche? *Folia Histochem. Cytobiol.* 45: 141-147.
5. Zheng, P. and Dean, J. 2007. Oocyte-specific genes affect folliculogenesis, fertilization, and early development. *Semin. Reprod. Med.* 25: 243-251.
6. Online Mendelian Inheritance in Man, OMIM™. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 610224. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
7. Choi, Y., Yuan, D. and Rajkovic, A. 2008. Germ cell-specific transcriptional regulator SOHLH2 is essential for early mouse folliculogenesis and oocyte-specific gene expression. *Biol. Reprod.* 79: 1176-1182.
8. Toyoda, S., Miyazaki, T., Miyazaki, S., Yoshimura, T., Yamamoto, M., Tashiro, F., Yamato, E. and Miyazaki, J. 2009. SOHLH2 affects differentiation of KIT positive oocytes and spermatogonia. *Dev. Biol.* 325: 238-248.

CHROMOSOMAL LOCATION

Genetic locus: SOHLH1 (human) mapping to 9q34.3.

PRODUCT

SOHLH1 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SOHLH1 shRNA Plasmid (h): sc-76537-SH and SOHLH1 shRNA (h) Lentiviral Particles: sc-76537-V as alternate gene silencing products.

For independent verification of SOHLH1 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-76537A, sc-76537B and sc-76537C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

SOHLH1 siRNA (h) is recommended for the inhibition of SOHLH1 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SOHLH1 gene expression knockdown using RT-PCR Primer: SOHLH1 (h)-PR: sc-76537-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.