

TNF α -IP 1 siRNA (m): sc-76697

BACKGROUND

TNF α -IP 1 (tumor necrosis factor, α -induced protein 1, endothelial), also known as B12, B61, EDP1 or TNFAIP1, is a BTB/POZ domain-containing protein that belongs to the KCTD10/KCTD13/TNFAIP1 family. TNF α -IP 1 contains one BTB/POZ motif, which is known to mediate homomeric and heteromeric POZ-POZ interactions and is common to transcriptional regulators involved in chromatin modeling. The expression of TNF α -IP 1 can be induced by IL-6 (interleukin-6) and by TNF α in the umbilical vein of endothelial cells. TNF α -IP 1 may be involved in DNA repair, DNA synthesis, cell apoptosis and human diseases. TNF α -IP 1 is suggested to play a role in the process of cancer and in the innate immunity against the Hepatitis B virus.

REFERENCES

1. Swift, S., et al. 1998. Structure and chromosomal mapping of the TNF α inducible endothelial protein 1 (Edp1) gene in the mouse. *Biochim. Biophys. Acta* 1442: 394-398.
2. Link, C.D., et al. 2003. Gene expression analysis in a transgenic *Caenorhabditis elegans* Alzheimer's disease model. *Neurobiol. Aging* 24: 397-413.
3. Zhou, J., et al. 2005. Cloning of two rat PDIP1 related genes and their interactions with proliferating cell nuclear antigen. *J. Exp. Zool. Part A Comp. Exp. Biol.* 303: 227-240.
4. Lin, M.C., et al. 2005. Tumor necrosis factor α -induced protein 1 and immunity to Hepatitis B virus. *World J. Gastroenterol.* 11: 7564-7568.
5. Yang, L.P., et al. 2006. Expression profile in the cell lines of human TNFAIP1 gene. *Yi Chuan* 28: 918-922.
6. Ghanim, H., et al. 2007. Role of inflammatory mediators in the suppression of Insulin receptor phosphorylation in circulating mononuclear cells of obese subjects. *Diabetologia* 50: 278-285.
7. Wang, M., et al. 2008. Comparative analysis of transcriptional profiling of CD3⁺, CD4⁺ and CD8⁺ T cells identifies novel immune response players in T cell activation. *BMC Genomics* 9: 225.

CHROMOSOMAL LOCATION

Genetic locus: Tnfaip1 (mouse) mapping to 11 B5.

PRODUCT

TNF α -IP 1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TNF α -IP 1 shRNA Plasmid (m): sc-76697-SH and TNF α -IP 1 shRNA (m) Lentiviral Particles: sc-76697-V as alternate gene silencing products.

For independent verification of TNF α -IP 1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-76697A, sc-76697B and sc-76697C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TNF α -IP 1 siRNA (m) is recommended for the inhibition of TNF α -IP 1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TNF α -IP 1 gene expression knockdown using RT-PCR Primer: TNF α -IP 1 (m)-PR: sc-76697-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Liu, N., et al. 2016. TNFAIP1 contributes to the neurotoxicity induced by A β ₂₅₋₃₅ in Neuro2a cells. *BMC Neurosci.* 17: 51.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.