

TRAC-1 siRNA (h): sc-76723

BACKGROUND

The RING-type zinc finger motif is present in a number of viral and eukaryotic proteins and is made of a conserved cysteine-rich domain that is able to bind two zinc atoms. Proteins that contain this conserved domain are generally involved in the ubiquitination pathway of protein degradation. TRAC-1 (T cell RING activation protein 1), also known as RNF125 (RING finger protein 125) or E3 ubiquitin-protein ligase RNF125, is a 232 amino acid novel E3 ubiquitin ligase that functions as a positive regulator in the T cell receptor signaling pathway. Expressed predominantly in lymphoid tissues such as spleen, thymus and bone marrow, TRAC-1 has been found to inhibit pathogen-induced cytokine production and downregulates HIV replication.

REFERENCES

- Saurin, A.J., et al. 1996. Does this have a familiar RING? *Trends Biochem. Sci.* 21: 208-214.
- Zhao, H., et al. 2005. A novel E3 ubiquitin ligase TRAC-1 positively regulates T cell activation. *J. Immunol.* 174: 5288-5297.
- Arimoto, K., et al. 2007. Negative regulation of the RIG-I signaling by the ubiquitin ligase RNF125. *Proc. Natl. Acad. Sci. USA* 104: 7500-7505.
- Shoji-Kawata, S., et al. 2007. The RING finger ubiquitin ligase RNF125/TRAC-1 down-modulates HIV-1 replication in primary human peripheral blood mononuclear cells. *Virology* 368: 191-204.
- Online Mendelian Inheritance in Man, OMIM[™]. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 610432. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Giannini, A.L., et al. 2008. T cell regulator RNF125/TRAC-1 belongs to a novel family of ubiquitin ligases with zinc fingers and a ubiquitin-binding domain. *Biochem. J.* 410: 101-111.
- Arimoto, K., et al. 2008. UbcH8 regulates ubiquitin and ISG15 conjugation to RIG-I. *Mol. Immunol.* 45: 1078-1084.

CHROMOSOMAL LOCATION

Genetic locus: RNF125 (human) mapping to 18q12.1.

PRODUCT

TRAC-1 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TRAC-1 shRNA Plasmid (h): sc-76723-SH and TRAC-1 shRNA (h) Lentiviral Particles: sc-76723-V as alternate gene silencing products.

For independent verification of TRAC-1 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-76723A, sc-76723B and sc-76723C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TRAC-1 siRNA (h) is recommended for the inhibition of TRAC-1 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TRAC-1 gene expression knockdown using RT-PCR Primer: TRAC-1 (h)-PR: sc-76723-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.