

UBE2Q2 siRNA (m): sc-76793

BACKGROUND

UBE2Q2 (ubiquitin-conjugating enzyme E2 Q2) is a 375 amino acid cytoplasmic protein that is involved in ubiquitin-mediated protein degradation. Ubiquitination is an important mechanism through which three classes of enzymes act in concert to target short-lived or abnormal proteins for destruction. The three classes of enzymes involved in ubiquitination are the ubiquitin-activating enzymes (E1s), the ubiquitin-conjugating enzymes (E2s) and the ubiquitin-protein ligases (E3s). One of several members of the ubiquitin-conjugating enzyme family, UBE2Q2 is an E2 ubiquitin-conjugating enzyme that acts to catalyze the covalent attachment of ubiquitin residues to various proteins. UBE2Q2 expression is detected in invasive epithelial tissue and tumor masses including head and neck squamous cell carcinomas, suggesting a role for UBE2Q2 in carcinogenesis. Two isoforms of UBE2Q2 exist due to alternative splicing events.

REFERENCES

- Schulze, E., Altmann, M.E., Adham, I.M., Schulze, B., Fröde, S. and Engel, W. 2003. The maintenance of neuromuscular function requires UBC-25 in *Caenorhabditis elegans*. *Biochem. Biophys. Res. Commun.* 305: 691-699.
- Melner, M.H., Ducharme, N.A., Brash, A.R., Winfrey, V.P. and Olson, G.E. 2004. Differential expression of genes in the endometrium at implantation: upregulation of a novel member of the E2 class of ubiquitin-conjugating enzymes. *Biol. Reprod.* 70: 406-414.
- Zhang, Y.W., Otterness, D.M., Chiang, G.G., Xie, W., Liu, Y.C., Mercurio, F. and Abraham, R.T. 2005. Genotoxic stress targets human Chk1 for degradation by the ubiquitin-proteasome pathway. *Mol. Cell* 19: 607-618.
- Melner, M.H., Haas, A.L., Klein, J.M., Brash, A.R., Boeglin, W.E., Nagdas, S.K., Winfrey, V.P. and Olson, G.E. 2006. Demonstration of ubiquitin thiolester formation of UBE2Q2 (UBCi), a novel ubiquitin-conjugating enzyme with implantation site-specific expression. *Biol. Reprod.* 75: 395-406.
- Seghatoleslam, A., Zambrano, A., Millon, R., Ganguli, G., Argentini, M., Cromer, A., Abecassis, J. and Wasylyk, B. 2006. Analysis of a novel human gene, LOC92912, over-expressed in hypopharyngeal tumours. *Biochem. Biophys. Res. Commun.* 339: 422-429.
- Banerjee, S., Brooks, W.S. and Crawford, D.F. 2007. Inactivation of the ubiquitin conjugating enzyme UBE2Q2 causes a prophase arrest and enhanced apoptosis in response to microtubule inhibiting agents. *Oncogene* 26: 6509-6517.

CHROMOSOMAL LOCATION

Genetic locus: Ube2q2 (mouse) mapping to 9 B.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

UBE2Q2 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see UBE2Q2 shRNA Plasmid (m): sc-76793-SH and UBE2Q2 shRNA (m) Lentiviral Particles: sc-76793-V as alternate gene silencing products.

For independent verification of UBE2Q2 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-76793A, sc-76793B and sc-76793C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

UBE2Q2 siRNA (m) is recommended for the inhibition of UBE2Q2 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor UBE2Q2 gene expression knockdown using RT-PCR Primer: UBE2Q2 (m)-PR: sc-76793-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.