

ZNF442 siRNA (h): sc-76986

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZNF442 is a 627 amino acid member of the Krüppel C₂H₂-type zinc-finger protein family. Localized to the nucleus, ZNF442 contains 16 C₂H₂-type zinc fingers and one KRAB domain. The gene encoding ZNF442 maps to chromosome 19p13.2. Chromosome 19, which makes up over 2% of human genomic DNA, is the genetic home for a number of immunoglobulin superfamily members, including the killer cell and leukocyte Ig-like receptors, a number of ICAMs, the CEACAM and PSG family, and Fc α receptors.

REFERENCES

1. Rooney, P.H., Boonsong, A., McFadyen, M.C., McLeod, H.L., Cassidy, J., Curran, S. and Murray, G.I. 2004. The candidate oncogene ZNF217 is frequently amplified in colon cancer. *J. Pathol.* 204: 282-288.
2. Huang, G., Krig, S., Kowbel, D., Xu, H., Hyun, B., Volik, S., Feuerstein, B., Mills, G.B., Stokoe, D., Yaswen, P. and Collins, C. 2005. ZNF217 suppresses cell death associated with chemotherapy and telomere dysfunction. *Hum. Mol. Genet.* 14: 3219-3225.
3. Sarraf, S., Tejada, R., Abawi, M., Oberst, M., Dennis, T., Simon, K.C. and Blacato, J. 2005. The human ovarian teratocarcinoma cell line PA-1 demonstrates a single translocation: analysis with fluorescence *in situ* hybridization, spectral karyotyping, and bacterial artificial chromosome microarray. *Cancer Genet. Cytogenet.* 161: 63-69.
4. Shimada, M., Imura, J., Kozaki, T., Fujimori, T., Asakawa, S., Shimizu, N. and Kawaguchi, R. 2005. Detection of Her2/Neu, c-Myc and ZNF217 gene amplification during breast cancer progression using fluorescence *in situ* hybridization. *Oncol. Rep.* 13: 633-641.
5. Zhong, M., Li, J., Ding, Y.Q. and Song, L.L. 2006. ZNF217 gene was detected in ovarian serous cystadenocarcinoma by fluorescence *in situ* hybridization. *Zhonghua Yi Xue Yi Chuan Xue Za Zhi* 23: 665-667.
6. Quinlan, K.G., Verger, A., Yaswen, P. and Crossley, M. 2007. Amplification of zinc finger gene 217 (ZNF217) and cancer: when good fingers go bad. *Biochim. Biophys. Acta* 1775: 333-340.
7. Li, P., Maines-Bandiera, S., Kuo, W.L., Guan, Y., Sun, Y., Hills, M., Huang, G., Collins, C.C., Leung, P.C., Gray, J.W. and Auersperg, N. 2007. Multiple roles of the candidate oncogene ZNF217 in ovarian epithelial neoplastic progression. *Int. J. Cancer* 120: 1863-1873.
8. Krig, S.R., Jin, V.X., Bieda, M.C., O'Geen, H., Yaswen, P., Green, R. and Farnham, P.J. 2007. Identification of genes directly regulated by the oncogene ZNF217 using chromatin immunoprecipitation (ChIP)-chip assays. *J. Biol. Chem.* 282: 9703-9712.
9. Cowger, J.J., Zhao, Q., Isovich, M. and Torchia, J. 2007. Biochemical characterization of the zinc-finger protein 217 transcriptional repressor complex: identification of a ZNF217 consensus recognition sequence. *Oncogene* 26: 3378-3386.

CHROMOSOMAL LOCATION

Genetic locus: ZNF442 (human) mapping to 19p13.2.

PRODUCT

ZNF442 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ZNF442 shRNA Plasmid (h): sc-76986-SH and ZNF442 shRNA (h) Lentiviral Particles: sc-76986-V as alternate gene silencing products.

For independent verification of ZNF442 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-76986A, sc-76986B and sc-76986C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ZNF442 siRNA (h) is recommended for the inhibition of ZNF442 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ZNF442 gene expression knockdown using RT-PCR Primer: ZNF442 (h)-PR: sc-76986-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.