

ZNF516 siRNA (h): sc-76992

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. As a member of the Krüppel C₂H₂-type zinc-finger protein family, ZNF516 (zinc-finger protein 516) is a 1,163 amino acid nuclear protein that contains ten C₂H₂-type zinc-fingers. The gene encoding ZNF516 maps to human chromosome 18q23, in a region that is frequently found to be affected in 18q deletion syndrome, a multiple-anomaly mental retardation syndrome that is associated with congenital aural atresia.

REFERENCES

1. Payre, F., et al. 1988. Finger proteins and DNA-specific recognition: distinct patterns of conserved amino acids suggest different evolutionary modes. *FEBS Lett.* 234: 245-250.
2. Rosenfeld, R., et al. 1993. Zinc-fingers: conserved properties that can distinguish between spurious and actual DNA-binding motifs. *J. Biomol. Struct. Dyn.* 11: 557-570.
3. Nagase, T., et al. 1996. Prediction of the coding sequences of unidentified human genes. VI. The coding sequences of 80 new genes (KIAA0201-KIAA0280) deduced by analysis of cDNA clones from cell line KG-1 and brain. *DNA Res.* 3: 321-329, 341-354.
4. Urrutia, R. 2003. KRAB-containing zinc-finger repressor proteins. *Genome Biol.* 4: 231.
5. Nuijten, I., et al. 2003. Congenital aural atresia in 18q deletion or de Grouchy syndrome. *Otol. Neurotol.* 24: 900-906.
6. Beausoleil, S.A., et al. 2004. Large-scale characterization of HeLa cell nuclear phosphoproteins. *Proc. Natl. Acad. Sci. USA* 101: 12130-12135.
7. Dostal, A., et al. 2006. Identification of 2.3-Mb gene locus for congenital aural atresia in 18q22.3 deletion: a case report analyzed by comparative genomic hybridization. *Otol. Neurotol.* 27: 427-432.
8. Riley, D.E., et al. 2007. Simple repeat evolution includes dramatic primary sequence changes that conserve folding potential. *Biochem. Biophys. Res. Commun.* 355: 619-625.

CHROMOSOMAL LOCATION

Genetic locus: ZNF516 (human) mapping to 18q23.

PRODUCT

ZNF516 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ZNF516 shRNA Plasmid (h): sc-76992-SH and ZNF516 shRNA (h) Lentiviral Particles: sc-76992-V as alternate gene silencing products.

For independent verification of ZNF516 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-76992A, sc-76992B and sc-76992C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ZNF516 siRNA (h) is recommended for the inhibition of ZNF516 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ZNF516 gene expression knockdown using RT-PCR Primer: ZNF516 (h)-PR: sc-76992-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.