

EXOSC7 siRNA (m): sc-77295

BACKGROUND

The exosome is a multisubunit complex of 3' to 5' exoribonucleases. It is involved in a variety of cellular processes and is responsible for degrading unstable mRNAs that contain AU-rich elements in their untranslated 3' region. EXOSC7 (exosome component 7), also known as p8, EAP1, RRP42 (ribosomal RNA-processing protein 42), Rrp42p or hRrp42p, is a component of the exosome multienzyme ribonuclease complex. It belongs to the RNase PH family and localizes to the nucleolus. EXOSC7 is one of the six RNase-PH domain subunits of the exosome. Together, these six subunits form a PNPase-like RING. EXOSC7 is required for the processing of the 7S pre-RNA.

REFERENCES

1. Nagase, T., et al. 1995. Prediction of the coding sequences of unidentified human genes. III. The coding sequences of 40 new genes (KIAA0081-KIAA0120) deduced by analysis of cDNA clones from human cell line KG-1. *DNA Res.* 2: 37-43.
2. Chen, C.Y., et al. 2001. AU binding proteins recruit the exosome to degrade ARE-containing mRNAs. *Cell* 107: 451-464.
3. Raijmakers, R., et al. 2002. Protein-protein interactions of hCsl4p with other human exosome subunits. *J. Mol. Biol.* 315: 809-818.
4. Brouwer, R., et al. 2002. Autoantibodies directed to novel components of the PM/ScI complex, the human exosome. *Arthritis Res.* 4: 134-138.
5. Raijmakers, R., et al. 2002. Protein-protein interactions between human exosome components support the assembly of RNase PH-type subunits into a six-membered PNPase-like RING. *J. Mol. Biol.* 323: 653-663.
6. Online Mendelian Inheritance in Man, OMIM[™]. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 606488. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
7. Anderson, J.R., et al. 2006. Sequence-specific RNA binding mediated by the RNase PH domain of components of the exosome. *RNA* 12: 1810-1816.
8. Walter, P., et al. 2006. Characterization of native and reconstituted exosome complexes from the hyperthermophilic archaeon *Sulfolobus solfataricus*. *Mol. Microbiol.* 62: 1076-1089.

CHROMOSOMAL LOCATION

Genetic locus: Exosc7 (mouse) mapping to 9 F4.

PRODUCT

EXOSC7 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see EXOSC7 shRNA Plasmid (m): sc-77295-SH and EXOSC7 shRNA (m) Lentiviral Particles: sc-77295-V as alternate gene silencing products.

For independent verification of EXOSC7 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-77295A, sc-77295B and sc-77295C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

EXOSC7 siRNA (m) is recommended for the inhibition of EXOSC7 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

EXOSC7 (A-1): sc-393686 is recommended as a control antibody for monitoring of EXOSC7 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor EXOSC7 gene expression knockdown using RT-PCR Primer: EXOSC7 (m)-PR: sc-77295-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.