

DNASE1L2 siRNA (h): sc-77320

BACKGROUND

DNASE1L2 (deoxyribonuclease I-like 2), also known as DHP1 or DNAS1L2, is a 299 amino acid secreted protein that is expressed in brain tissue and shares sequence similarity with DNase I, suggesting a possibly role in DNA hydrolysis. The gene encoding DNASE1L2 maps to human chromosome 16, which encodes over 900 genes and comprises nearly 3% of the human genome. The GAN gene is located on chromosome 16 and, with mutation, may lead to giant axonal neuropathy, a nervous system disorder characterized by increasing malfunction with growth. The rare disorder Rubinstein-Taybi syndrome is also associated with chromosome 16, as is Crohn's disease, which is a gastrointestinal inflammatory condition.

REFERENCES

- Rodriguez, A.M., Rodin, D., Nomura, H., Morton, C.C., Weremowicz, S. and Schneider, M.C. 1997. Identification, localization, and expression of two novel human genes similar to deoxyribonuclease I. *Genomics* 42: 507-513.
- Shiokawa, D. and Tanuma, S. 2001. Characterization of human DNase I family endonucleases and activation of DNase γ during apoptosis. *Biochemistry* 40: 143-152.
- Shiokawa, D., Matsushita, T., Kobayashi, T., Matsumoto, Y. and Tanuma, S. 2004. Characterization of the human DNAS1L2 gene and the molecular mechanism for its transcriptional activation induced by inflammatory cytokines. *Genomics* 84: 95-105.
- Coupry, I., Monnet, L., Attia, A.A., Taine, L., Lacombe, D. and Arveiler, B. 2004. Analysis of CBP (CREBBP) gene deletions in Rubinstein-Taybi syndrome patients using real-time quantitative PCR. *Hum. Mutat.* 23: 278-284.
- Online Mendelian Inheritance in Man, OMIM[™]. 2004. Johns Hopkins University, Baltimore, MD. MIM Number: 602622. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Jager, K., Fischer, H., Tschachler, E. and Eckhart, L. 2007. Terminal differentiation of nail matrix keratinocytes involves up-regulation of DNASE1L2 but is independent of caspase-14 expression. *Differentiation* 75: 939-946.
- Fischer, H., Eckhart, L., Mildner, M., Jaeger, K., Buchberger, M., Ghannadan, M. and Tschachler, E. 2007. DNASE1L2 degrades nuclear DNA during corneocyte formation. *J. Invest. Dermatol.* 127: 24-30.

CHROMOSOMAL LOCATION

Genetic locus: DNASE1L2 (human) mapping to 16p13.3.

PRODUCT

DNASE1L2 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see DNASE1L2 shRNA Plasmid (h): sc-77320-SH and DNASE1L2 shRNA (h) Lentiviral Particles: sc-77320-V as alternate gene silencing products.

For independent verification of DNASE1L2 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-77320A, sc-77320B and sc-77320C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

DNASE1L2 siRNA (h) is recommended for the inhibition of DNASE1L2 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor DNASE1L2 gene expression knockdown using RT-PCR Primer: DNASE1L2 (h)-PR: sc-77320-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.