



TMEM70 siRNA (h): sc-77756

BACKGROUND

TMEM70 (transmembrane protein 70) is a 260 amino acid multi-pass mitochondrial membrane protein that exists as multiple alternatively spliced isoforms which are expressed in pancreas, liver, muscle and kidney tissue. Mutations in the gene encoding TMEM70 are associated with ATP synthase deficiency and neonatal mitochondrial encephalomyopathy, suggesting that TMEM70 plays an important role in ADP-stimulated respiration, mitochondrial ATP synthesis, ATP hydrolysis and the maintenance of membrane potential. The gene encoding TMEM70 maps to human chromosome 8, which consists of nearly 146 million base pairs, houses more than 800 genes and is associated with a variety of diseases and malignancies.

REFERENCES

1. Wildenauer, D.B. and Schwab, S.G. 1999. Chromosomes 8 and 10 workshop. *Am. J. Med. Genet.* 88: 239-243.
2. Calvo, S., Jain, M., Xie, X., Sheth, S.A., Chang, B., Goldberger, O.A., Spinazzola, A., Zeviani, M., Carr, S.A. and Mootha, V.K. 2006. Systematic identification of human mitochondrial disease genes through integrative genomics. *Nat. Genet.* 38: 576-582.
3. Cízková, A., Stránecký, V., Mayr, J.A., Tesarová, M., Havlíčková, V., Paul, J., Ivánek, R., Kuss, A.W., Hansíková, H., Kaplanová, V., Vrbáček, M., Hartmannová, H., Nosková, L., Honzíček, T., et al. 2008. TMEM70 mutations cause isolated ATP synthase deficiency and neonatal mitochondrial encephalomyopathy. *Nat. Genet.* 40: 1288-1290.
4. Online Mendelian Inheritance in Man, OMIM™. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 612418. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
5. Houstěk, J., Knoch, S. and Zeman, J. 2009. TMEM70 protein—a novel ancillary factor of mammalian ATP synthase. *Biochim. Biophys. Acta* 1787: 529-532.
6. Wortmann, S.B., Rodenburg, R.J., Jonckheere, A., de Vries, M.C., Huizing, M., Heldt, K., van den Heuvel, L.P., Wendel, U., Kluijtmans, L.A., Engelke, U.F., Wevers, R.A., Smeitink, J.A. and Morava, E. 2009. Biochemical and genetic analysis of 3-methylglutaconic aciduria type IV: a diagnostic strategy. *Brain* 132: 136-146.

CHROMOSOMAL LOCATION

Genetic locus: TMEM70 (human) mapping to 8q21.11.

PRODUCT

TMEM70 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TMEM70 shRNA Plasmid (h): sc-77756-SH and TMEM70 shRNA (h) Lentiviral Particles: sc-77756-V as alternate gene silencing products.

For independent verification of TMEM70 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-77756A, sc-77756B and sc-77756C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TMEM70 siRNA (h) is recommended for the inhibition of TMEM70 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

TMEM70 (B-6): sc-393619 is recommended as a control antibody for monitoring of TMEM70 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TMEM70 gene expression knockdown using RT-PCR Primer: TMEM70 (h)-PR: sc-77756-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.